

**Max-Planck-Institut
für Mathematik
in den Naturwissenschaften
Leipzig**

**Unexpected solutions of first and second
order partial differential equations**

by

Stefan Müller and Vladimír Šverák

Preprint-Nr.: 42

1998



Unexpected solutions of first and second order partial differential equations

S. Müller ¹

Max Planck Institute for
Mathematics in the Sciences
Inselstr. 22-24, D-04103 Leipzig

and

V. Šverák

School of Mathematics
University of Minnesota
206 Church Street SE
Minneapolis, MN 55455

1 Introduction and examples

In this paper we discuss a general approach to construct solutions for a large class of nonlinear first and second order partial differential equations. The method makes strong use of work by Gromov (who substantially generalized earlier results of Nash and Kuiper) and is especially suitable for nonconvex problems where standard compactness arguments fail. One application concerns the (unexpected) existence of solutions in mathematical models of solid-solid phase transformations (see Example c) below). Another application is the recent resolution of the regularity question for weak solutions to the Euler-Lagrange equations of multiple integrals.

Theorem 1.1 *There exists a smooth, strongly elliptic 2×2 system*

$$-\operatorname{div} \sigma(Dv) = 0, \quad v : \mathbb{R}^2 \rightarrow \mathbb{R}^2 \quad (1.1)$$

that admits

- (i) nontrivial Lipschitz solutions with compact support;*
- (ii) Lipschitz solutions that are nowhere C^1 .*

Moreover σ can be chosen such that (1.1) is the Euler-Lagrange equation of a variational integral $\int f(Dv) dx$, where f is smooth and uniformly quasiconvex in the sense of Morrey.

The existence of irregular solutions of the Euler-Lagrange equations (1.1) is in sharp contrast with the (partial) regularity theory for minimizers of quasiconvex integrals ([Ev 86], [AF 87], [GM 86], [FH 85]).

¹The authors were supported by the Max Planck research award

This raises the question which structure conditions on σ are needed to ensure a good regularity theory for quasilinear systems. (Tartar has raised this issue in connection with the closely related question of compactness and stability of solutions, see e.g. [Ta 79], p. 160.) For a scalar equation the De Giorgi-Moser-Nash Theorem shows that ellipticity is the natural condition. For systems, there is a large literature for monotone σ (see [Gi 83] for a summary, further references and a sketch of the history) and many results can be extended to so-called quasimonotone σ ([Fu 87], [Ha 95], [Zh 88]), but these conditions are too restrictive for applications e.g. to nonlinear elasticity. Similar issues arise for problems in nondivergence form as recent counterexamples by Nadi-rashvili show ([Na 98]). In the theory of harmonic maps there are also striking differences between minimizers, weak solutions of the Euler-Lagrange equations and so-called stationary harmonic maps which in addition satisfy a form of the Euler-Lagrange equations obtained by variation of the independent variable ([Mo 48], [SU 82], [He 91], [Ev 91], [Be 93], [Ri 95]; see [He 97], [Si 96] for recent overviews).

We remark in passing that our counterexamples to regularity are quite different from the famous examples of De Giorgi [DG 68], Giusti-Miranda [GM 68] Bombieri-De Giorgi-Giusti [BDG 69] and many subsequent works including [HLN 96]. The latter are based on finding equations that admit certain point singularities like $x/|x|$ (or certain cones), while our approach uses the fact that the equation is compatible with certain large oscillations of Du (small oscillations must be smooth by ellipticity). The construction of counterexamples is thus reduced to certain algebraic calculations in the space of matrices (see Section 4 below). Our point of view here is strongly influenced by the work of Tartar [Ta 79], [Ta 83], [Ta 98] and by the result [Sv 95] that there exists a 6×2 system that admits a smooth periodic solution (and arises as the Euler-Lagrange equation of a strictly polyconvex variational integral).

Before we return to the case of 2×2 systems let us review the general setting and some illustrative examples. Given a subset of the $m \times n$ matrices $M^{m \times n}$, a (bounded) domain $\Omega \subset \mathbb{R}^n$ and a map $u_0 : \Omega \rightarrow \mathbb{R}^m$ we seek to find Lipschitz maps $u : \Omega \rightarrow \mathbb{R}^m$ that satisfy

$$Du(x) \in K \quad \text{for a.e. } x \in \Omega, \quad (1.2)$$

$$u = u_0 \quad \text{on } \partial\Omega. \quad (1.3)$$

Generalizations to problems of the form $F(x, u(x), Du(x)) = 0$ a.e., to maps between manifolds and to higher order derivatives are possible. In order to avoid technicalities as much as possible we focus on (1.2) and (1.3) in the following. This setting already includes a number of interesting examples.

Example a) (scalar u , Hamilton-Jacobi equations) Let $m = 1$. It follows from Theorem 2.4 below that (1.2), (1.3) has a solution if u_0 is C^1 (or piecewise C^1 and continuous) and

$$Du_0 \in \text{int conv} K.$$

For affine u_0 the condition $Du_0 \in \overline{\text{conv}} K$ is clearly necessary. On the other hand, the examples $K = \{a, b\}$ or $K = \{(\pm 1, \pm 1)\}$ show that the condition $Du_0 \in \overline{\text{conv}} K$ is in general not sufficient, even if u_0 is affine. As a special case

consider a convex, coercive function $f : \mathbb{R}^n \rightarrow \mathbb{R}$. Then the Hamilton-Jacobi equation

$$f(Du) = 0 \tag{1.4}$$

has a solution which satisfies the boundary condition (1.3) provided that

$$f(Du_0) < 0$$

In this case there is also a good theory of viscosity solutions of (1.4) (see [Kr 75], [CL 83]). In general, Theorem 2.4 below yields existence of solutions in many cases where no viscosity solution exists, but the solutions have much weaker properties (no uniqueness, no comparison principle), see [CDGG 99] for further discussion.

b) (Isometries) If $K = SO(n)$ then Liouville's theorem shows that Du is constant on each component of Ω (see [Re 68] and [BI 82], or [Ki 88] for the simpler Lipschitz case). On the other hand if $K = O(n)$ or

$$K = O(n, m) = \{F \in M^{m \times n} : F^T F = id_{\mathbb{R}^n}\}$$

then (1.2), (1.3) admits a solution if u_0 is a 'short' map, i.e.

$$Du_0 \in \text{int conv} K = \{F \in M^{m \times n} : \lambda_{max}(F^T F) < 1\},$$

see [Gr 86], Chapter 2.4.11, p. 216, [CP 95], [CP 97]. In fact for $m > n$ one can obtain C^1 solutions [Gr 86], Chapter 2.4.9, Thm. (A), p. 203.

c) (Two-well problem) In the study of phase transitions in crystals ([BJ 87], [BJ 92], [Bh 92], [CK 88]; see [Mu 98] for a recent survey) the set

$$K = SO(2)A \cup SO(2)B \subset M^{2 \times 2},$$

with A, B symmetric, positive definite, $\det A = \det B = 1$ arises. Theorem 3.2. below shows that solutions exist if $u_0 \in C^{1, \alpha}$ (for $0 < \alpha < 1$) and

$$Du_0 \in \text{int } \overline{\text{conv}} K \cap \{\det = 1\}.$$

d) ($m \times 2$ elliptic systems) Let $\sigma : M^{m \times 2} \rightarrow M^{m \times 2}$ be a C^1 map and consider the second order system

$$-\text{div } \sigma(Dv) = 0 \quad \text{in } \Omega, \tag{1.5}$$

i.e.

$$-\sum_{\alpha=1}^2 \partial_{\alpha} \sigma_{i\alpha}(Dv) = 0, \quad \text{for } i = 1, \dots, m.$$

If Ω is simply connected then (1.5) can be expressed as

$$\sigma(Dv)J = Dw, \quad J = \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix},$$

and if we let $u = \begin{pmatrix} v \\ w \end{pmatrix}$, then (1.5) can be rewritten as

$$Du \in K, \quad K = \left\{ \begin{pmatrix} F \\ G \end{pmatrix} \in M^{2m \times 2} : \sigma(F)J = G \right\},$$

and thus can be recast into the present framework.

e) (Four-point configuration). The following example played an important rôle in clarifying different convexity notions in the calculus of variations and was discovered independently (in different contexts) by several authors ([AH 86], [CT 93], [Ta 93]; for a related example see [MN 91]). It will be crucial in the construction of nontrivial solutions to 2×2 elliptic systems. Let (see Figure 1 in section 4)

$$K = \{A_1, A_2, A_3, A_4\}, \quad -A_1 = A_3 = \begin{pmatrix} 1 & 0 \\ 0 & 3 \end{pmatrix}, \quad -A_2 = A_4 = \begin{pmatrix} 3 & 0 \\ 0 & -1 \end{pmatrix}.$$

One easily checks that all solutions of $Du \in K$ are trivial. Corollary 4.1 below shows that there is a large number of nontrivial maps whose gradient stays in an arbitrarily small neighbourhood of K .

2 Convex integration

The first striking results on solutions of relations like (1.2) appeared in the fundamental work of Nash [Na 54] and Kuiper [Ku 55] on isometric immersions. Specifically, Kuiper showed that for any $\varepsilon > 0$ there exist an isometric C^1 immersion $u : S^2 \rightarrow \mathbb{R}^3$ that maps S^2 in a ball of radius ε , while a classical theorem of Hilbert states that C^2 isometric immersions are rigid motions (see Borisov [Bo 65] for rigidity and non-rigidity in $C^{1,\alpha}$). Extending these ideas Gromov [Gr 73, Gr 86] developed a general method, called ‘convex integration’ to treat (1.2) and much more general partial differential relations (Spring’s recent book [Sp 98] gives a detailed exposition). The main emphasis in [Gr 86] is on the construction of C^1 solutions. In the context of equidimensional isometric immersions Gromov also studies the Lipschitz case in detail and later states a general result for Lipschitz solutions, see Chapter 2.4.11, p. 218. The setting is that of jet bundles and thus the result covers in particular systems of the form $F(x, u(x), \dots, D^{(m)}u(x)) = 0$ a.e. in Ω subject to $D^{(l)}u = v^{(l)}$ on $\partial\Omega$, $0 \leq l \leq m-1$.

A short self-contained proof for the special case (1.2), (1.3) appeared in [MS 96]. A slightly different approach based on Baire’s theorem was pursued by Dacorogna and Marcellini [DM 97], [DM 98a], [DM 98b], following work of Cellina [Ce 80] as well as De Blasi and Pianigiani [DP 82, DP 91] for ordinary differential equation and differential inclusions. As we shall see Gromov’s setting (or that of Dacorogna and Marcellini) suffices to discuss Examples a) and b), while for c)–e) additional ideas are needed.

The basic idea of convex integration is that nontrivial solutions of (1.2), (1.3) exist if Du_0 takes values in (the interior of) a suitable convex hull, called the P -convex hull. For sets $K \subset M^{m \times n}$ the notion of P -convexity reduces to what is called lamination convexity in [MS 96] ([MP 98] use the term set-theoretic rank-1 convexity). A set $E \subset M^{m \times n}$ is lamination convex if for all matrices $A, B \in E$ that satisfy $\text{rk}(B - A) = 1$, the whole segment $[A, B]$ is contained in E .

The lamination convex hull E^{lc} is the smallest lamination convex set containing E . The relevance of rank-1 matrices stems from the fact that they arise exactly as gradients of maps $x \mapsto u(x \cdot n)$ which only depends on one variable. These maps (and slight modifications thereof; see Lemma 2.2 below) are the basic building blocks in Gromov's construction. In the scalar case $m = 1$ lamination convexity of course reduces to ordinary convexity.

The construction of solutions now proceeds in two steps. First one considers open sets $U \subset M^{m \times n}$, and this case is easily reduced to an open neighbourhood of two matrices A, B with $\text{rk}(B - A) = 1$. Secondly one passes to general sets K by approximating them from the inside by open sets contained in K^{lc} . In the following we say that a map $u : \Omega \subset \mathbb{R}^n \rightarrow \mathbb{R}^m$ is piecewise linear if it is continuous, if there exist finitely or countably many disjoint sets Ω_i whose union has full measure such that $u|_{\Omega_i}$ is affine and if Du is (essentially) bounded.

Lemma 2.1 *Suppose that $U \subset M^{m \times n}$ is open and bounded and that $u_0 : \Omega \rightarrow \mathbb{R}^m$ is piecewise linear and satisfies*

$$Du_0 \in U^{lc} \text{ a.e.} \quad (2.1)$$

Then there exists, for any $\delta > 0$, a piecewise linear u such that

$$Du \in U \text{ a.e., } u = u_0 \text{ on } \partial\Omega, \quad (2.2)$$

$$\sup |u - u_0| < \delta. \quad (2.3)$$

For the proof it clearly suffices to consider the case $u_0(x) = Fx$. One uses the fact that U^{lc} can be defined by successive addition of rank-1 segments:

$$\begin{aligned} U^{lc} &= \bigcup_{k=0}^{\infty} U^{(k)}, \text{ where } U^{(0)} = U, \\ U^{(k+1)} &= U^{(k)} \cup \{[A, B] : A, B \in U, \text{rk}(B - A) = 1\}. \end{aligned}$$

By induction over k the proof of Lemma 2.1 is easily reduced to the following special case (see [MS 96]).

Lemma 2.2 *Suppose that $\text{rk}(B - A) = 1$, i.e. $B - A = a \otimes n$, and let $F = \lambda A + (1 - \lambda)B$. If U is an open neighbourhood of $\{A, B\}$ then there exists a piecewise linear u such that*

$$Du \in U \text{ a.e., } u = Fx \text{ on } \partial\Omega, \quad (2.4)$$

$$\sup |u(x) - Fx| < \delta. \quad (2.5)$$

To construct u , assume without loss of generality $F = 0, n = e_n$ and consider the set $M = (-1, 1)^{n-1} \times \varepsilon(-\lambda, 1 - \lambda)$, the one-dimensional map $v(x) = ah(x_n)$, where $h'(x_n) = 1 - \lambda$ for $x_n < 0$, $h'(x_n) = -\lambda$ for $x_n > 0$, $h(0) = \varepsilon\lambda(1 - \lambda)$. Then $Dv \in \{A, B\}$ and $h > 0$ in M . The function $u(x) = ag(x)$, with $g(x) = h(x) - \varepsilon \sum_{i=1}^{n-1} |x_i|$ has the desired properties on the diamond-shaped set $\tilde{M} = M \cap \{g > 0\}$. For general sets $\Omega \subset \mathbb{R}^n$ one can use Vitali's theorem to exhaust Ω by disjoint scaled copies of \tilde{M} . Choosing the scaled copies sufficiently small one obtains (2.5).

This finishes the argument for open sets U . For general sets K one needs an suitable approximation by open sets.

Definition 2.3 (Gromov) *A sequence of open sets $U_i \subset M^{m \times n}$ is an in-approximation of a set $E \subset M^{m \times n}$ if*

- (i) *the U_i are uniformly bounded;*
- (ii) *$U_i \subset U_{i+1}^{lc}$;*
- (iii) *$U_i \rightarrow E$ in the following sense: if $F_i \in U_i$ and $F_i \rightarrow F$ then $F \in E$.*

Example For $m = 1$ the shells $U_i = \{x : 1 - 2^{-i+2} < |x| < 1\}$ are an in-approximation of S^{n-1} .

Theorem 2.4 ([Gr 86, p. 218; [MS 96]]) *Suppose that $\{U_i\}$ is an in-approximation of $K \subset M^{m \times n}$ and that $u_0 : \Omega \rightarrow \mathbb{R}^m$ is C^1 (or piecewise C^1) and satisfies*

$$Du_0 \in U_1 \text{ a.e.}$$

Then there exists $u \in W^{1,\infty}(\Omega; \mathbb{R}^m)$ such that

$$Du \in K \text{ a.e., } u = u_0 \text{ on } \partial\Omega$$

For the proof one uses Lemma 2.1 to inductively construct approximations $u^{(i)}$ with $Du^{(i)} \in U_i$. The key point is to assure that the $Du^{(i)}$ converge strongly. At first glance it is surprising that this can be achieved since the construction in Lemma 2.1 yields solutions with highly oscillatory gradients. Nonetheless by a judicious choice of the C^0 error δ in Lemma 2.1 one can ensure that the oscillations added in each iteration step are essentially independent of the previous ones and only effect a set of small measure. This construction, which is reminiscent of the construction of continuous, nowhere differentiable functions is one of the key ideas of convex integration (in [DM 97] it is replaced by an elegant, but slightly less flexible, Baire category argument). Specifically, one choses $\varepsilon_i < \frac{1}{2}$ and δ_i inductively such that $\|\rho_{\varepsilon_i} * Du_i - Du_i\|_{L^1} < 2^{-i}$ and $\sup |u_i - u_{i+1}| < \delta_{i+1} := \varepsilon_i \delta_i$, where $\rho_\varepsilon = \varepsilon^{-n} \rho(\frac{\cdot}{\varepsilon})$ is the usual mollifier. Then $u_i \rightarrow u_\infty$ in C^0 and $Du_i \rightarrow Du_\infty$ in L^1 since $\|\rho_{\varepsilon_i} * (Du_i - Du_\infty)\|_{L^1} = \|D\rho_{\varepsilon_i} * (u_i - u_\infty)\|_{L^1} \leq \frac{C\delta_{i+1}}{\varepsilon_i} \rightarrow 0$; see [MS 96] for the details.

3 Constraints and sets without rank-1 connections

The theory explained so far applies to Examples a) and b) but not to c) - e). As regards c), the constraint $\det F = 1$ is stable under lamination convexity since $F \mapsto \det F$ is affine in rank-1 directions. Hence the set K in c) does not admit an in-approximation by open sets. The set K in e) contains no rank-1 connections and hence $K^{lc} = K$ and similarly U^{lc} contains only points near K for small neighbourhoods U of K . As regards d), Ball [Ba 80] showed that for strongly elliptic systems that arise as Euler-Lagrange equations (i.e. $\sigma = Df, f : M^{m \times n} \rightarrow \mathbb{R}$) again $K^{lc} = K$. It turns out, however, that the previous results can be extended to a slightly larger hull than K^{lc} , namely the rank-1 convex hull K^{rc} (called the functional rank-1 convex hull in [MP 98]) and that this hull can be nontrivial in Examples d) and e).

We say that a function $f : E \rightarrow \mathbb{R}$ is rank-one convex on a set E if it is convex on each rank-one line $t \mapsto F + ta \otimes n$. For a compact set K we define the rank-one convex hull relative to E by

$$K^{rc,E} = \left\{ F \in M^{m \times n} : f(F) \leq \inf_K f, \forall f : E \rightarrow \mathbb{R} \text{ rank-1 convex} \right\},$$

i.e. $K^{rc,E}$ consists of those points that cannot be separated from K by rank-1 convex functions. For an open set U the set $U^{rc,E}$ is defined as the union of all $K^{rc,E}$ where $K \subset U$ is compact (some authors alternatively define the hull of an open set as $\overline{U}^{rc,E}$ but the above definition is more convenient and in line with the usual convex hull). If $E = M^{m \times n}$ we simply write K^{rc} and U^{rc} . The main result is the following variant of Lemma 2.1. Given an $r \times r$ minor M ($r \geq 2$) and a real number $t \neq 0$ we let

$$\Sigma = \{ F \in M^{m \times n} : M(F) = t \}.$$

Lemma 3.1 (i) *Let $U \subset M^{m \times n}$ be open, let $F \in U^{rc}$ and let $\varepsilon > 0$. Then there exists a piecewise linear map $u : \Omega \rightarrow \mathbb{R}^m$ that satisfies*

$$\begin{aligned} Du &\in U^{rc} \text{ a.e., } u(x) = Fx \text{ on } \partial\Omega, \\ \text{meas } \{ Du \notin U \} &< \varepsilon \text{ meas } \Omega. \end{aligned}$$

(ii) *If U is relatively open in Σ and $F \in U^{rc,\Sigma}$, then u can be chosen such that in addition $Du \in U^{rc,\Sigma} \subset \Sigma$ a.e.*

By a simple iteration one obtains the counterpart of Lemma 2.1 with U^{lc} replaced by U^{rc} (or $U^{rc,E}$ if a constraint is imposed). The proof of part(i) uses three facts. First, for a compact set K , the rank-1 convex hull K^{rc} consists of the barycentres of a certain class $\mathcal{M}^{rc}(K)$ ('laminates') of probability measures supported on K . Precisely, a probability measure belongs to $\mathcal{M}^{rc}(K)$ if and only if $\langle \nu, f \rangle \geq f(\langle \nu, id \rangle)$ for all rank-1 convex f . Secondly, we use a result of Pedregal [Pe 93] that laminates can be approximated (in the weak* topology of measures) by simpler measures, the so-called laminates of finite order, that are supported on U^{rc} , where U is a (small) neighbourhood of K . The class $\mathcal{L}(U^{rc})$ of laminates of finite order is defined inductively as follows: all Dirac masses δ_F with $F \in U^{rc}$ belong to $\mathcal{L}(U^{rc})$. If $\sum_{i=1}^k \lambda_i \delta_{F_i}$ belongs to $\mathcal{L}(U^{rc})$ and if $F_k = \mu A + (1 - \mu)B$, $A, B \in U^{rc}$, $\text{rk}(B - A) = 1$, $\mu \in (0, 1)$, then $\sum_{i=1}^{k-1} \lambda_i \delta_{F_i} + \lambda_k \mu \delta_A + \lambda_k (1 - \mu) \delta_B$ belongs to $\mathcal{L}(U^{rc})$. Third, we inductively use Lemma 2.2 to associate to each $\nu \in \mathcal{L}(U^{rc})$ a map $u : \Omega \rightarrow \mathbb{R}^m$ such that $Du \in U^{rc}$ and $|\text{meas } \{ Du = F_i \} - \lambda_i \text{ meas } \Omega| < 2^{-i} \varepsilon$. Then u has the desired properties.

To treat the case with constraint one first has to extend Pedregal's result to this situation. Secondly one has to prove a version of Lemma 2.2 which includes the constraint $Du \in \Sigma$ (one can relax (2.4)₁ to the conditions $Du \in U^{rc}$ and $\text{meas } \{ Du \notin U \} < \varepsilon$.) To this end one first obtains a C^∞ approximation by smoothing and considering a flow of a suitable (divergence-free) vectorfield. Then one constructs a piecewise linear approximation using, among other facts, a result of Dacorogna and Moser [DM 90] on the solvability of $\det Du = f$ in $C^{k,\alpha}$ spaces.

By iteration and the same approximation argument as in the proof of Theorem 2.4 one finally obtains the following result. We say that $\{U_i\}$ is an *rc-in-approximation* of K if the conditions in Definition 2.3 hold with U_i^{lc} replaced by U_i^{rc} .

Theorem 3.2 (i) Suppose that $K \in M^{m \times n}$ admits an *rc-in-approximation* by open sets U_i . Suppose that $u_0 \in C^1(\Omega; \mathbb{R}^m)$ (or u_0 piecewise C^1) and

$$Du_0 \in U_1^{rc}.$$

Then there exists a map u that satisfies

$$Du \in K \text{ a.e., } u = u_0 \text{ on } \partial\Omega.$$

(ii) If $K \subset \Sigma$ and $u_0 \in C^{2,\alpha}(\Omega; \mathbb{R}^m)$ then the same assertion holds if the U_i are only relatively open in Σ . If $m = n = r$ then the condition $u_0 \in C^{1,\alpha}(\Omega; \mathbb{R}^m)$ suffices.

Remark. If K is open (or relatively open in Σ) one can take the trivial in-approximation $U_i \equiv K$.

4 Applications

Example a) Lamination convexity reduces to ordinary convexity, and an in-approximation with $U_1 \supset \text{int conv } K$ is given by $U_i = \{F \in \text{conv } K : 0 < \text{dist}(F, K) < 2^{-i+2} \text{diam } K\}$. Hence the assertions in Section 1 follow from Theorem 2.4.

Example b) One easily checks that

$$K^{lc} = K^{rc} = \text{conv } K = \{F \in M^{m \times n} : \lambda_{\max}(F^T F) \leq 1\}.$$

It follows that

$$U_i = \{F \in M^{m \times n} : 1 - 2^{-i+2} < \lambda_{\max}(F^T F) < 1\}$$

provides an in-approximation with $U_1 = \text{int conv } K$, and Theorem 2.4 applies.

Example c) Let $\Sigma = \{F \in M^{2 \times 2} : \det F = 1\}$. By a result of Šverák [Sv 93a]

$$K^{lc} = K^{rc} = \Sigma \cap \text{conv } K.$$

Thus $U_i = \Sigma \cap \{F \in \text{int conv } K : 0 < \text{dist}(F, K) < 2^{-i+2} \text{diam } K\}$ provides an in-approximation (relative to Σ) with $U_1 \supset \Sigma \cap \text{int conv } K$.

Example e) Let

$$J_1 = -J_3 = \begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix}, \quad J_2 = -J_4 = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}.$$

Then $K^{lc} = K$ but

$$K^{rc} = [-1, 1]^2 \cup \bigcup_{i=1}^4 [J_i, A_{i+1}]$$

(see Figure 1).

As an immediate consequence of Theorem 3.2 we obtain:




Figure 1: The set $\{A_1, A_2, A_3, A_4\}$ is lamination convex but the rank-one convex hull contains the shaded square and the line segments $[J_i, A_{i+1}]$.

Corollary 4.1 *Let $U \supset K$ be open, and suppose that $F \in U^{rc} \supset K^{rc}$. Then there exist $u : \Omega \rightarrow \mathbb{R}^2$ such that*

$$Du \in U \text{ a.e., } u = Fx \text{ on } \partial\Omega.$$

As regards Example d) we recall the result mentioned in the introduction.

Theorem 4.2 ([MS 98b]) *There exists a smooth strongly elliptic 2×2 system*

$$-\operatorname{div} \sigma(Dv) = 0, \quad v : \mathbb{R}^2 \rightarrow \mathbb{R}^2 \quad (4.1)$$

that admits

- (i) *nontrivial Lipschitz solutions with compact support;*
- (ii) *Lipschitz solutions that are nowhere C^1 .*

Moreover σ can be chosen such that (4.1) is the Euler-Lagrange equation of a variational integral $\int f(Dv) dx$, where f is smooth and uniformly quasiconvex in the sense of Morrey.

Sketch of proof. Our interest lies mainly in the variational case but the main idea can already be seen in the simpler non-variational situation. The key idea is to embed the four-point configuration in Figure 1 in the set

$$K = \left\{ \begin{pmatrix} F \\ G \end{pmatrix} : F, G \in M^{2 \times 2}, \sigma(F)J = G \right\}.$$

This turns out to be surprisingly simple. Consider first the restriction of K to diagonal matrices and let

$$\begin{aligned}\sigma_{11}(F_{11}, F_{22}) &= F_{11} - g(F_{22}), \\ \sigma_{22}(F_{11}, F_{22}) &= F_{22} - h(F_{11}).\end{aligned}$$

Strong ellipticity on diagonal matrices reduces to the conditions

$$\frac{\partial \sigma_{11}}{\partial F_{11}} \geq c > 0, \quad \frac{\partial \sigma_{22}}{\partial F_{22}} \geq c > 0,$$

and is clearly satisfied. Moreover g and h can be chosen such that the set $\{\sigma_{11} = \sigma_{22} = 0\}$ includes the points A_1, A_2, A_3, A_4 in Figure 1 and $(0, 0)$. If we extend σ to nondiagonal matrices by

$$\sigma_{12} = kF_{12}, \quad \sigma_{21} = kF_{21}$$

then σ is elliptic for sufficiently large k , and a careful analysis shows that generically K^{rc} contains a neighbourhood U of $0 \in M^{4 \times 2}$, and K admits an rc-in-approximation $\{U_i\}$ with $U_1 = U$.

Let Ω be a smooth and bounded domain in \mathbb{R}^n . By Theorem 3.2 there exists a solution

$$Du \in K \text{ a.e.}, \quad u = 0 \text{ on } \partial\Omega.$$

Writing $u = \begin{pmatrix} v \\ w \end{pmatrix}$ we obtain

$$\begin{aligned}-\operatorname{div} \sigma(Dv) &= 0 \quad \text{in } \Omega, \\ v &= 0 \quad \text{on } \partial\Omega.\end{aligned}$$

Since $Dw = \sigma(Dv)J$, the trace theorem yields

$$\sigma(Dv)n = 0 \quad \text{on } \partial\Omega.$$

Now extend v by zero to \mathbb{R}^n . Since $\sigma(0) = 0$ the map v is a solution of (4.1) with compact support. Regarding (ii) one can use (i) and scaling to construct solutions that can only be regular on a set of arbitrarily small measure. To obtain the full strength of (ii) one has to slightly modify the construction in the proof of Theorem 2.4.

5 Some open problems

A necessary condition for the solvability of

$$Du \in K \text{ a.e. in } \Omega, \quad u(x) = Fx \text{ on } \partial\Omega$$

is that F belongs to the so-called quasiconvex hull K^{qc} of K which in general is bigger than the rank-1 convex hull K^{rc} (see [Sv 95] or [Mu 98] for definitions and further references). This raises the following questions

- Does Theorem 3.2 hold if one replaces U_i^{lc} by U_i^{qc} in the definition of in-approximation?

- Can one compute (or estimate) K^{qc} for the set K in Example d)?
- Can one find manageable conditions on σ that guarantee $K^{qc} = K$?

Even checking whether $K^{rc} = K$ is in general not obvious. The following Theorem gives a recent example.

Theorem 5.1 ([MS 98c]) *Let $f(F) = \det F$, for $F \in M^{2 \times 2}$, and let $\sigma(F) = Df(F) = \det F \operatorname{cof} F$. Then the set*

$$K = \left\{ \begin{pmatrix} F \\ G \end{pmatrix} : \sigma(F)J = G \right\}$$

satisfies $K^{rc} = K$.

Acknowledgements

The work reported here is the result of a long collaboration which was supported by the Alexander von Humboldt foundation through the Max Planck research award. We also greatly benefited from discussions with L. Tartar and B. Kirchheim. The IMA kindly provided office space and logistic support during several visits of SM at the University of Minnesota.

References

- [AF 87] E. Acerbi and N. Fusco, A regularity theorem for minimizers of quasi-convex integrals, *Arch. Rat. Mech. Anal.* **89** (1987), 251 – 281.
- [AH 86] R. Aumann and S. Hart, Bi-convexity and bi-martingales, *Israel J. Math.* **54** (1986), 159 – 180.
- [Ba 80] J.M. Ball, Strict convexity, strong ellipticity and regularity in the calculus of variations, *Math. Proc. Cambridge Phil. Soc.* **87** (1980), 501 – 513.
- [BJ 87] J.M. Ball and R.D. James, Fine phase mixtures as minimizers of energy, *Arch. Rat. Mech. Anal.* **100** (1987), 13 – 52.
- [BJ 92] J.M. Ball and R.D. James, Proposed experimental tests of a theory of fine microstructure and the two-well problem, *Phil. Trans. Roy. Soc. London A* **338** (1992), 389 – 450.
- [Be 93] F. Bethuel, On the singularity set of stationary harmonic maps, *manuscripta math.* **78** (1993), 417 – 443.
- [Bh 92] K. Bhattacharya, Self-accommodation in martensite, *Arch. Rat. Mech. Anal.* **120** (1992), 201 – 244.
- [BI 82] B.V. Bojarski and T. Iwaniec, Another approach to the Liouville theorem, *Math. Nachr.* **107** (1982), 253 – 262.
- [BDG 69] E. Bombieri, E. De Giorgi and E. Giusti, Minimal cones and the Bernstein theorem, *Invent. Math.* **7** (1969), 243–269.

- [Bo 65] Ju. Borisov $C^{1,\alpha}$ isometric immersions of Riemannian spaces, *Dokl. Akad. S.S.S.R* **163** (1965), 11 – 13.
- [CDGG 99] P. Cardaliaguet, B. Dacorogna, W. Gangbo, N. Georgy, Geometric restrictions for the existence of viscosity solutions, *Ann. IHP Analyse non linéaire*, to appear (1999).
- [CP 97] P. Celada and S. Perrotta, Functions with prescribed singular values of the gradient, preprint SISSA 56/97/M, 1997.
- [CT 93] E. Casadio-Tarabusi, An algebraic characterization of quasi-convex functions, *Ricerche Mat.* **42** (1993), 11 – 24.
- [Ce 80] A. Cellina, On the differential inclusion $x \in [-1, 1]$, *Atti Accad. Naz. Lincei Rend. Sci. Fis. Mat. Nat.* **69**, 1 – 6.
- [CP 95] A. Cellina and S. Perrotta, On a problem of potential wells, *J. convex anal.* **2** (1995), 103 – 115.
- [CK 88] M. Chipot and D. Kinderlehrer, Equilibrium configurations of crystals, *Arch. Rat. Mech. Anal.* **103** (1988), 237 – 277.
- [CL 83] M.G. Crandall and P.-L. Lions, Viscosity solutions of Hamilton-Jacobi equations, *Trans. Amer. Math. Soc.* **277** (1983), 1 – 42.
- [DM 97] B. Dacorogna and P. Marcellini, General existence theorems for Hamilton-Jacobi equations in the scalar and vectorial cases, *Acta Math.* **178** (1997), 1 – 37.
- [DM 98a] B. Dacorogna and P. Marcellini, Cauchy-Dirichlet problem for first order nonlinear systems, *J. Funct. Anal.* **152** (1998), 404 – 446.
- [DM 98b] B. Dacorogna and P. Marcellini, Implicit second order partial differential equations, *Ann. SNS Pisa*, to appear.
- [DM 90] B. Dacorogna and J. Moser, On a partial differential equation involving the Jacobian determinant, *Ann. IHP Analyse non linéaire* **7** (1990), 1–26.
- [DP 82] F.S. De Blasi and G. Pianigiani, Baire category approach to the existence of solutions of multivalued differential equations in Banach spaces, *Funkcial. Ekvac.* **25** (1982), 153–162.
- [DP 91] F.S. De Blasi and G. Pianigiani, Non convex valued differential inclusions in Banach spaces, *J. Math. Anal. Appl.* **157** (1991), 469–494.
- [DG 68] E. De Giorgi, Un esempio di estremali discontinue per un problema variazionale di tipo ellittico, *Boll. U.M.I.* **4** (1968), 135–137.
- [Ev 86] L.C. Evans, Quasiconvexity and partial regularity in the calculus of variations, *Arch. Rat. Mech. Anal.* **95** (1986), 227–252.
- [Ev 91] L.C. Evans, Partial regularity for stationary harmonic maps into spheres, *Arch. Rat. Mech. Anal.* **116**, 101 – 113.

- [Fu 87] M. Fuchs, Regularity theorems for nonlinear systems of partial differential equations under natural ellipticity conditions, *Analysis* **7** (1987), 83 – 93.
- [FH 85] N. Fusco and J. Hutchinson, Partial regularity of functions minimizing quasiconvex integrals, *manuscripta math.* **54** (1985), 121–143.
- [Gi 83] M. Giaquinta, *Multiple integrals in the calculus of variations and nonlinear elliptic systems*, Princeton UP, 1983.
- [GM 86] M. Giaquinta and G. Modica, Partial regularity of minimizers of quasiconvex integrals, *Ann. IHP Analyse non linéaire* **3** (1986), 185 – 208.
- [GM 68] E. Giusti and M. Miranda, Un esempio di soluzioni discontinue per un problema di minimo relativo ad un integrale regolare del calcolo delle variazioni, *Boll. U.M.I* **2** (1968), 1–8.
- [Gr 73] M. Gromov, Convex integration of differential relations, *Izw. Akad. Nauk S.S.S.R* **37** (1973), 329 – 343.
- [Gr 86] M. Gromov, *Partial differential relations*, Springer, 1986.
- [Ha 95] C. Hamburger, Quasimonotonicity, regularity and duality for nonlinear systems of partial differential equations, *Ann. Mat. Pure Appl. Ser. IV* **169** (1995), 321 – 354.
- [HLN 96] W. Hao, S. Leonardi, J. Nečas An example of irregular solution to a nonlinear Euler-Lagrange elliptic system with real analytic coefficients. *Ann. SNS Pisa Ser. IV* **23** (1996), 57–67.
- [He 91] F. Hélein, Régularité des applications faiblement harmoniques entre une surface et une variété Riemannienne, *C.R.A.S. Paris* **312** (1991), 591 – 596.
- [He 97] F. Hélein, *Harmonic maps, conservation laws and moving frames*, Diderot, 1997.
- [Ki 88] D. Kinderlehrer, Remarks about equilibrium configurations of crystals, in: *Material instabilities in continuum mechanics and related mathematical problems* (J.M. Ball, ed.), Oxford Univ. Press, 1988, 217 – 242.
- [Kr 75] S.N. Kruzkov, Generalized solutions of Hamilton-Jacobi equation of eikonal type, *USSR Sbornik* **27** (1975), 406 – 446.
- [Ku 55] N.H. Kuiper, On C^1 isometric embeddings, I., *Nederl. Akad. Wetensch. Proc. A* **58** (1955), 545 – 556.
- [MP 98] J. Matoušek and P. Plecháč, On functional separately convex hulls, *J. Discrete Comput. Geom.* **19** (1998), 105 – 130.
- [Mi 98] G.W. Milton, *The effective tensors of composites*, book in preparation.
- [MN 91] V. Nesi and G.W. Milton, Polycrystalline configurations that maximize electrical resistivity, *J. Mech. Phys. Solids* **39** (1991), 525–542.
- [Mo 48] C.B. Morrey, The problem of Plateau on a Riemannian manifold, *Ann. Math.* **49** (1948), 807 – 851.

- [Mo 66] C.B. Morrey, *Multiple integrals in the calculus of variations*, Springer, 1966.
- [MS 96] S. Müller and V. Šverák, Attainment results for the two-well problem by convex integration, *Geometric analysis and the calculus of variations*, (J. Jost, ed.), International Press, 1996, 239 – 251.
- [Mu 98] S. Müller, Variational models for microstructure and phase transitions, to appear in: Proc. C.I.M.E summer school ‘Calculus of variations and geometric evolution problems’ (S. Hildebrandt and M. Struwe, eds.), <http://www.mis.mpg.de/cgi-bin/lecturenotes.pl>.
- [MS 98a] S. Müller and V. Šverák, Convex integration with constraints and applications to phase transitions and partial differential equations, in preparation.
- [MS 98b] S. Müller and V. Šverák, in preparation.
- [MS 98c] S. Müller and V. Šverák, in preparation.
- [Na 98] N. Nadirashvili, Nonuniqueness in the martingale problem and the Dirichlet problem for uniformly elliptic operators, *Ann. SNS Pisa Ser. IV* **24** (1997), 537 – 550.
- [Na 54] J. Nash, C^1 isometric embeddings, *Ann. Math.* **60** (1954), 383–396.
- [Pe 93] P. Pedregal, Laminates and microstructure, *Europ. J. Appl. Math.* **4** (1993), 121–149.
- [Re 68] Yu.G. Reshetnyak, Liouville’s theorem under minimal regularity assumptions, *Sib. Math. J.* **8** (1968), 631–634.
- [Ri 95] T. Rivière, Everywhere discontinuous harmonic maps into spheres, *Acta Math.* **175** (1995), 197 – 226.
- [SU 82] R. Schoen and K. Uhlenbeck, A regularity theory for harmonic maps, *J. Diff. Geom.* **17** (1982), 302 – 336.
- [Si 96] L. Simon, *Theorems on regularity and singularity of energy minimizing maps*, Lectures in Mathematics ETH Zürich, Birkhäuser, 1996.
- [Sp 98] D. Spring, *Convex integration theory*, Birkhäuser, 1998.
- [Sv 92] V. Šverák, Rank-one convexity does not imply quasiconvexity, *Proc. Roy. Soc. Edinburgh* **120** (1992), 185–189.
- [Sv 93a] V. Šverák, On the problem of two wells, in: *Microstructure and phase transitions*, IMA Vol. Appl. Math. **54** (D. Kinderlehrer, R.D. James, M. Luskin and J. Ericksen, eds.), Springer, 1993, 183 – 189.
- [Sv 95] V. Šverák, Lower semicontinuity of variational integrals and compensated compactness, in: Proc. ICM 1994 (S.D. Chatterji, ed.), vol. 2, Birkhäuser, 1995, 1153–1158.

- [Ta 79] L. Tartar, Compensated compactness and partial differential equations, in: *Nonlinear Analysis and Mechanics: Heriot-Watt Symposium* Vol. IV, (R. Knops, ed.), Pitman, 1979, 136–212.
- [Ta 83] L. Tartar, The compensated compactness method applied to systems of conservations laws, in: *Systems of Nonlinear Partial Differential Equations*, (J.M. Ball, ed.), NATO ASI Series, Vol. C111, Reidel, 1983, 263–285.
- [Ta 93] L. Tartar, Some remarks on separately convex functions, in: *Microstructure and phase transitions*, IMA Vol. Math. Appl. **54**, (D. Kinderlehrer, R.D. James, M. Luskin and J.L. Ericksen, eds.), Springer, 1993, 191–204.
- [Ta 98] L. Tartar, *Homogenization, compensated compactness and H-measures*, CBMS-NSF conference, Santa Cruz, June 1993, lecture notes in preparation.
- [Zh 88] Kewei Zhang, On the Dirichlet problem for a class of quasilinear elliptic systems of partial differential equations in divergence form, in: *Partial differential equations* (Tianjun, 1986), Lecture Notes in Mathematics **1306**, Springer, 1988.