

Max-Planck-Institut
für Mathematik
in den Naturwissenschaften
Leipzig

Global Existence for a Nonlinear
System in Thermoviscoelasticity with
Nonconvex Energy

by

Johannes Zimmer

Preprint no.: 100

2003



Global Existence for a Nonlinear System in Thermoviscoelasticity with Nonconvex Energy

Johannes Zimmer

*Max Planck Institute for Mathematics in the Sciences,
Inselstr. 22, 04 103 Leipzig, Germany
E-mail: zimmer@mis.mpg.de*

A three-dimensional thermoviscoelastic system derived from the balance laws of momentum and energy is considered. To describe structural phase transitions in solids, the stored energy function is not assumed to be convex as a function of the deformation gradient. A novel feature for multi-dimensional, nonconvex, and non-isothermal problems is that no regularizing higher order terms are introduced. The mechanical dissipation is not linearized. We prove existence global in time. The approach is based on a fixed-point argument using an implicit time discretization and the theory of renormalized solutions for parabolic equations with L^1 data.

1. INTRODUCTION

This article is concerned with global solvability of an initial-boundary value problem in three-dimensional thermoviscoelasticity arising from the theory of solid-solid phase transitions. We consider a three-dimensional body, identified with its reference configuration Ω in \mathbb{R}^3 . Here, Ω is assumed to be a bounded, nonempty domain with Lipschitz boundary. Let $T > 0$ be an arbitrary, but fixed time. The thermomechanical evolution of the body will be described in terms of the deformation field $u: \Omega \times [0, T[\rightarrow \mathbb{R}^n$ and the temperature field $\theta: \Omega \times [0, T[\rightarrow \mathbb{R}$. The evolution of the body also depends on the stored energy function $\Phi(F, \theta): \text{Mat}(n \times n) \times \mathbb{R} \rightarrow \mathbb{R}$ which enters the equation through the stress tensor $\sigma(F, \theta) = \frac{\partial \Phi(F, \theta)}{\partial F}$.

The balance laws of momentum and energy ultimately lead to the nonlinear coupled system

$$u_{tt} = \text{Div}(\sigma(\nabla u, \theta) + \nabla u_t) \quad \text{in } \Omega \times]0, T[, \quad (1a)$$

$$\theta_t = \Delta \theta + \theta \sigma_\theta(\nabla u, \theta) : \nabla u_t + \nabla u_t : \nabla u_t \quad \text{in } \Omega \times]0, T[. \quad (1b)$$

Initial and boundary conditions are specified in Section 2. Remarks about the derivation of this system follow at the end of this section.

To model phase transitions, we do *not* assume σ to be monotone in F for temperatures below the critical temperature (i.e., the temperature at which the phase transition occurs). Hence, the stored energy Φ will be nonconvex as a function of F below the critical temperature.

Some remarks about the peculiarities of this model and related systems are in order. From now on, we will focus on systems with nonconvex energies. Equations of thermoviscoelasticity with convex energy have been considered in [2], where renormalized solutions are used to show existence of a solution. We will rely heavily on this machinery and comment later on differences between convex and nonconvex energies. The novelty of the existence result presented in this paper (Theorem 3.1) is that, to our knowledge, all previous results for similar systems with nonconvex energies either

- (i) study the one-dimensional case, i.e., $\Omega = [0, 1]$,

or

- (ii) in the multi-dimensional case, concentrate on the isothermal problem or include capillarity-like higher-order terms that have a regularizing effect.

Even under these assumptions, it has been remarkably difficult to prove global existence. Let us highlight a few results.

- The one-dimensional case has been studied for a long time. See, e.g., [6, 7, 17, 5]. Recent advances have been made in [22], where rather weak solutions are considered. In the remarkable paper [26], Watson studies solids as well as gaseous materials. It is important to notice that all these results do not carry over to the multi-dimensional case: though very different techniques are used throughout these papers, the crucial step is always a bound of u_x in $L^\infty(\Omega; \mathbb{R})$. In higher space dimensions, even in the isothermal case, this estimate is *wrong* unless one imposes rotational invariance of Ω or physically unrealistic smoothness assumptions on the given data [24].

- The three-dimensional isothermal case has been studied by Rybka [23]. Among others, he proves existence and uniqueness of a solution in the multi-dimensional isothermal case for Lipschitz continuous σ . Later, this result has been generalized by Friesecke and Dolzmann [14] to allow a more general kind of nonlinearity in σ . Our proof uses the ideas developed in [14]. The main difficulty to generalize their results to the non-isothermal case is the mechanical dissipation $\nabla u_t : \nabla u_t$ in the heat equation.

- The three-dimensional, non-isothermal case including capillarity-like higher-order terms has been an open problem for some years until recently Pałwó and Żochowski [19] proved global existence. The proof relies on the

observation that the higher-order term allows a parabolic decomposition of the equation of motion. Preliminary results were, for example, obtained in [15].

System (1a)–(1b) is derived from the the balance laws of momentum and energy,

$$u_{tt} = \operatorname{Div} (\sigma(\nabla u, \theta) + \nabla u_t) \quad \text{in } \Omega \times]0, T[, \quad (2a)$$

$$\begin{aligned} -\theta \Phi_{\theta\theta}(\nabla u, \theta) \theta_t &= \Delta \theta + \theta \sigma_{\theta}(\nabla u, \theta) : \nabla u_t + \nabla u_t : \nabla u_t \\ &\quad \text{in } \Omega \times]0, T[. \end{aligned} \quad (2b)$$

In Section 2, the assumptions on Φ are listed. In particular, by (E2), one has

$$-\theta \Phi_{\theta\theta}(\nabla u, \theta) = 1 - \theta \phi''(\theta) \Phi_1(F).$$

For physical reasons, one expects $\phi(\theta)$ to be linear in θ . In this case, $1 - \theta \phi''(\theta) \Phi_1(F) = 1$, which means that (2a)–(2b) reduces to the system (1a)–(1b) under consideration. Yet, the growth conditions we must impose allow Φ to be linear only in an arbitrarily large, but fixed range of θ , so (1a)–(1b) represent the balance of energy within this restricted range. This explains why we replace the nonlinear term $1 - \theta \Phi_{\theta\theta}(\nabla u, \theta)$ by 1 and consequently study the system (1a)–(1b) as an approximation of the full system (2a)–(2b).

It should be noted that the viscous part ∇u_t of the stress tensor is not frame-indifferent. This is a typical weakness of multidimensional models of thermoviscoelasticity that are analytically tractable.

We will use a self-explanatory notation for Lebesgue and Sobolev space. E.g., $L^1(\Omega; \mathbb{R}^n)$ denotes the class of Lebesgue integrable functions defined on Ω with values in \mathbb{R}^n . Time-dependent function spaces will be denoted, e.g., $W^{1,2}(0, T; W^{1,p}(\Omega; \mathbb{R}))$.

To prove global existence of a solution to the system (1a)–(1b), we combine methods developed in [14] (to deal with the nonconvexity) with the theory of renormalized solutions in classical thermoviscoelasticity (to deal with the mechanical dissipation; see in particular the paper by Blanchard and Guibé [2]).

2. THE INITIAL-BOUNDARY VALUE PROBLEM

The system (1a)–(1b) has to be furnished with appropriate initial and boundary conditions. We study the initial-boundary value problem

$$u_{tt} = \operatorname{Div} (\sigma(\nabla u, \theta) + \nabla u_t) \quad \text{in } \Omega \times]0, T[, \quad (3a)$$

$$\begin{aligned} \theta_t &= \Delta\theta + \theta\sigma_\theta(\nabla u, \theta) : \nabla u_t + \nabla u_t : \nabla u_t \\ &\text{in } \Omega \times]0, T[, \end{aligned} \quad (3b)$$

$$u = g \quad \text{on } \partial\Omega \times [0, T[, \quad (3c)$$

$$u = u_0 \quad \text{in } \Omega \times \{0\}, \quad (3d)$$

$$u_t = v_0 \quad \text{in } \Omega \times \{0\}, \quad (3e)$$

$$\theta = 0 \quad \text{on } \partial\Omega \times [0, T[, \quad (3f)$$

$$\theta = \theta_0 \quad \text{in } \Omega \times \{0\}. \quad (3g)$$

Here, g, u_0, v_0 and θ_0 are given functions. Their regularity is specified in Section 2.1. The boundary condition $\theta = 0$ on $\partial\Omega$ seems to be inappropriate, since this corresponds to zero (absolute) temperature on the boundary. However, after reformulating the above equations for the incremental temperature field with respect to a fixed temperature $\bar{\Theta}$, rather than for the absolute temperature θ , one can see that the same proof holds for (positive) boundary data $\bar{\Theta}$. We refrain from spelling this out, to keep the notation simple.

On the stored energy $\Phi(F, \theta)$, the following conditions will be imposed:

1. Φ is sufficiently smooth:

$$\Phi \in C^2(\text{Mat}(n \times n) \times \mathbb{R}; \mathbb{R}). \quad (E1)$$

2. The stored energy is of the form

$$\Phi(F, \theta) = \alpha + \theta - \theta \ln(\theta) + \phi(\theta)\Phi_1(F) + \Phi_2(F), \quad (E2)$$

where α is constant. The other quantities will be specified in the next paragraphs.

To simplify the notation, let us define

$$\phi_j(F) := \frac{\partial \Phi_j(F)}{\partial F} \quad (j = 1, 2). \quad (4)$$

3. Growth condition on $\phi: \mathbb{R} \rightarrow \mathbb{R}$:

$$|\phi(\theta)|, |\theta\phi'(\theta)|, |\phi'(\theta)| \leq C. \quad (E3)$$

4. Growth condition on Φ_1 :

$$\Phi_1 \text{ has linear growth near infinity. That is, } \phi_1 \text{ is bounded.} \quad (E4)$$

Additionally, we require that

$$\phi_1 \text{ is a globally Lipschitz continuous function.} \quad (E5)$$

5. Growth condition on Φ_2 :

$$\phi_2 = \frac{\partial \Phi_2}{\partial F} \text{ is a globally Lipschitz continuous function,} \quad (\text{E6})$$

and there exist $c, c' > 0$ such that

$$c|F|^2 - c' \leq \Phi_2(F) \leq c'(|F|^2 + 1) \text{ and } |\phi_2(F)| \leq c'(|F| + 1). \quad (\text{E7})$$

Remark 2. 1. We need to explain that the growth conditions (E1)–(E7) are consistent with experimental observations of martensitic phase transitions. First, it is an inherent property of martensitic phase transitions that the energy is convex for large temperature (beyond the so-called M_d temperature, about 300°C for NiTi). And as there are no nonconvexities at high temperature, there are no nonconvexities for large strains. The reason is that only the parent (austenitic) and the martensitic phases are stable, and their strains differ only by a few percent. Phases with large strain are always unstable, which means they are in the convex region of the energy landscape, away from the minimizers. Therefore, for large temperatures or strains, the problem reduces to one which is very similar to the one studied by Blanchard and Guibé [2]. The challenge is exactly to address the nonconvexity in small strains and below the transition temperature. Indeed, it is important to notice that even the modeling implicitly relies on the small strain assumption, which makes the difficult mathematical question of growth conditions at Infinity largely irrelevant from a point of view of applications. Specifically, the modeling of phase transitions on the continuum level relies on the existence of Ericksen-Pitteri neighborhoods [12, 20]. This approach is only valid if the different stable phases can be confined to a neighborhood which does, on the crystalline level, not include shifts by one atomic layer (the cut-off function ξ in the example below singles out an analogue neighborhood on the continuum level). The fact that martensitic phase transitions are a small strain phenomenon is illustrated in [1]. It is demonstrated there that for specific phase transitions like the fcc-to-bcc transition occurring in iron, the crystalline energy necessarily has to be in L^∞ . Finally, we remark that some of the conditions stated above are familiar from related problems: the conditions imposed on the temperature in (E3) for $\phi(\theta)$ can also be found, for example, in [25] (however, the restrictive assumption (E4) is not imposed there).

Remark 2. 2. A priori, it is not even clear that the strong growth conditions one has to impose on the energy to obtain existence results of multi-dimensional thermoviscoelastic models of phase transitions (even with higher-order terms [19]) can be met for a frame-indifferent function. However, a general method to derive energy functions meeting all physical




FIG. 1. Left: Prototypical shape of $\phi(\theta)$, satisfying (E3). Right: The cut-off function ξ used in the example in Remark 2.2. It takes the value 1 in a region surrounding the possibly stable phases, and decreases to 0 for large strains.

requirements (and arbitrary growth conditions) is presented in [27, 28]. For illustrational purposes, we give an example of the tetragonal-orthorhombic (orthoI) symmetry breaking, as it occurs in Zirconia (ZrO_2). As explained in [11], this problem can be studied in two space dimensions. To ensure frame-indifference of the energy (that is, to satisfy $\Phi(QF) = \Phi(F)$ for every $Q \in \text{SO}(3)$), we use the polar decomposition and write the energy in $C := F^T F$. We will write both $\Phi = \Phi(F)$ and $\Phi = \Phi(C)$ when no confusion can arise. It is convenient to introduce the Voigt notation for the components of C ,

$$C = \begin{pmatrix} c_1 & \frac{1}{2}c_6 \\ \frac{1}{2}c_6 & c_2 \end{pmatrix}.$$

Using the ideas of [27, 28, 11], one can easily see that every function in C with tetragonal symmetry can be written as

$$\Phi(c_1, c_2, c_6) := \bar{\Phi}(c_1 + c_2, c_1^2 + c_2^2, c_6^2).$$

We derive an energy function for a first order phase transition. Let us introduce $\rho_1(c_1, c_2, c_6) := c_1 + c_2$ and $\rho_2(c_1, c_2, c_6) := c_1^2 + c_2^2$. Then, if $\xi = \xi(c_1, c_2, c_6)$ is a smooth cut-off function (see Figure 1 on the right) and $\gamma > 0$ is a constant, a possible energy is given by

$$\begin{aligned} \Phi(c_1, c_2, c_3) &:= \alpha + \theta - \theta \ln(\theta) \\ &+ \sum_{j=1}^2 \xi \left[\arctan \left(\theta - \theta_c + \frac{1}{4} \right) \cdot \rho_j(c_1, c_2, c_6) \right. \\ &\quad \left. - \frac{1}{2} \rho_j(c_1, c_2, c_6)^2 + \frac{1}{3} (c_1, c_2, c_6)^3 \right] \\ &+ \xi \cdot c_6^2 + \gamma(c_1 + c_2). \end{aligned} \tag{5}$$

We first motivate this structure of Φ . The right-hand side of the first line of (5) gives the caloric term. The second and third line of (5) describe the behavior for ‘small’ strains, including the strain region in which the phase

transition occurs. The last line models the growth rate for ‘large’ strains in c_1, c_2 and the behavior in the off-diagonal c_6 , which is independent of the phase transition. The function in square brackets is the usual Landau-Ginzburg energy for first order transitions [13], where the planar symmetry has been factored out and the commonly used term $\theta - \theta_c + \frac{1}{4}$ has been replaced by $\arctan(\theta - \theta_c + \frac{1}{4})$. The elimination of the the planar symmetry is a physical necessity; one avoids unwanted minimizers. The growth conditions (E3) on θ are introduced for mathematical reasons. Obviously, Φ is of the type (E2) and satisfies the smoothness assumption (E1). It is easy to see that $\phi(\theta) := \arctan(\theta - \theta_c + \frac{1}{4})$ satisfies (E3), and $\Phi_2(F) := \xi c_6^2 + \gamma(c_1 + c_2)$ satisfies for large strains $\Phi_2(F) = \gamma(c_1 + c_2) = \gamma|F|^2$. Consequently, the growth conditions (E6), (E7) are met. The example shows that these growth conditions are physically reasonable (an experimental determination of the growth condition for large strains is not possible; in engineering literature, the use of piecewise quadratic functions is common [16]). Finally, the two remaining growth conditions (E4) and (E5) hold since we use the cut-off function ξ in the definition of Φ . Though these growth conditions are introduced for mathematical convenience, the discussion in the preceding remark provides a physical justification. Namely, (E4) and (E5) model the phase transition, which happens only in a bounded set of strains. The cut-off function ξ singles out such a set. Since we just want to give a prototypical example for an energy function, we refrain from introducing appropriate constants and parameters. In practice, one had to choose γ small enough to make sure Φ_2 does not hide the potential wells, and one had to choose ξ in such a way that its decrease does not destroy the convexity away from the minimizers. Both can easily be accomplished. The details are explained in [11], where it is also shown how to fit the location of minimizers, the energy barriers, and the elastic moduli. Since the result is necessarily more technical and does not yield deeper insight for our goals, we refrain from repeating it here.

2.1. Initial and Boundary Conditions

The given data should satisfy the following smoothness assumptions:

$$\begin{aligned} g &\in W^{1,2}(\Omega; \mathbb{R}^n), \\ u_0 &\in W_g^{1,2}(\Omega; \mathbb{R}^n) := \{u \in W^{1,2}(\Omega; \mathbb{R}^n) \mid u - g \in W_0^{1,2}(\Omega; \mathbb{R}^n)\}, \\ v_0 &\in L^2(\Omega; \mathbb{R}^n), \\ \theta_0 &\in L^1(\Omega; \mathbb{R}). \end{aligned}$$

2.2. Resulting System of Equations

Using (E2), one obtains

$$\sigma(F, \theta) = \frac{\partial \Phi(F, \theta)}{\partial F} = \phi(\theta)\phi_1(F) + \phi_2(F)$$

and $\theta\sigma_\theta(\nabla u, \theta) : \nabla u_t = \theta\phi'(\theta)\phi_1(\nabla u) : \nabla u_t$. To simplify the notation, let us write $f(\theta) := \theta\phi'(\theta)$. Then, system (3a)–(3g) becomes

$$u_{tt} = \text{Div}(\phi(\theta)\phi_1(\nabla u) + \phi_2(\nabla u) + \nabla u_t) \quad \text{in } \Omega \times]0, T[, \quad (6a)$$

$$u = g \quad \text{on } \partial\Omega \times [0, T[, \quad (6b)$$

$$u = u_0 \quad \text{in } \Omega \times \{0\}, \quad (6c)$$

$$u_t = v_0 \quad \text{in } \Omega \times \{0\}, \quad (6d)$$

$$\theta_t = \Delta\theta + f(\theta)\phi_1(\nabla u) : \nabla u_t + \nabla u_t : \nabla u_t \quad \text{in } \Omega \times]0, T[, \quad (7a)$$

$$\theta = 0 \quad \text{on } \partial\Omega \times [0, T[, \quad (7b)$$

$$\theta = \theta_0 \quad \text{in } \Omega \times \{0\}. \quad (7c)$$

3. EXISTENCE OF A WEAK-RENORMALIZED SOLUTION GLOBAL IN TIME

In order to prove existence of a weak solution, we have to overcome two main difficulties: the nonconvexity of the energy density and the mechanical dissipation, i.e., the term $\nabla u_t : \nabla u_t$ in the heat equation.

The latter will lead to a parabolic equation with initial data in $L^1(\Omega; \mathbb{R})$ and a right-hand side in $L^1(0, T; L^1(\Omega; \mathbb{R}))$. We will use the concept of renormalized solutions, introduced by Lions and DiPerna in their investigation of the Boltzmann equation [10, 9]. Further references for renormalized solutions for parabolic equations in L^1 are, e.g., [18, 4, 3]. Other frameworks are, for example, SOLA [8], and entropy solutions [21].

In this paper, we combine ideas of [14] and [2]. In the latter paper, renormalized solutions are applied to a thermoviscoelastic system with a convex stored energy. Apart from the free energy, the coupling of the equations studied here differs from [2]: for phase transitions, the essential parameter is the temperature θ , not its gradient $\nabla\theta$.

3.1. Quick Review of Renormalized Solutions

For the reader's convenience, the basic properties of renormalized solutions are briefly recollected. With our application in mind, we concentrate on parabolic equations.

This Section deals with equations of the type

$$\theta_t - \Delta\theta = H \text{ in } \Omega \times]0, T[, \quad (8a)$$

$$\theta = 0 \text{ on } \partial\Omega \times]0, T[, \quad (8b)$$

$$\theta = \theta_0 \text{ in } \Omega \times \{0\}, \quad (8c)$$

with $H \in L^1(0, T; L^1(\Omega; \mathbb{R}))$ and $\theta_0 \in L^1(\Omega; \mathbb{R})$.

$T_K(r) := \max(\min(r, K), -K)$ denotes the truncation function at height $K \geq 0$.

DEFINITION 3.1. A measurable function $\theta: \Omega \times]0, T[\rightarrow \mathbb{R}$ is a *renormalized solution* of problem (8a)–(8c) if it satisfies the following properties:

- (i) $\theta \in L^\infty(0, T; L^1(\Omega; \mathbb{R}))$,
- (ii) $T_K(\theta) \in L^2(0, T; W_0^{1,2}(\Omega; \mathbb{R}))$ for every $K \geq 0$,
- (iii) $\lim_{n \rightarrow +\infty} \iint_{\{(x,t) \mid n \leq |\theta(x,t)| \leq n+1\}} |\nabla\theta|^2 \, dy \, ds = 0$,

and, for every $S \in C^\infty(\mathbb{R}; \mathbb{R})$ with $S' \in C_0^\infty(\mathbb{R}; \mathbb{R})$,

$$(iv) \quad S(\theta)_t - \operatorname{div}[S'(\theta)\nabla\theta] + S''(\theta)|\nabla\theta|^2 = HS'(\theta) \quad \text{in } \mathcal{D}'(\Omega \times]0, T[),$$

$$(v) \quad S(\theta) = S(\theta_0) \text{ in } \Omega \times \{0\}.$$

Remark 3. 1. In our application, the right hand side H of the heat equation will depend on a temperature $\hat{\theta}$ and u :

$$\theta_t - \Delta\theta = H(\hat{\theta}, u) \text{ in } \Omega \times]0, T[,$$

where $H(\hat{\theta}, u) := f(\hat{\theta})\phi_1(\nabla u) : \nabla u_t + \nabla u_t : \nabla u_t$. To obtain the regularity $H \in L^1(0, T; L^1(\Omega; \mathbb{R}))$, seeking for $u \in W^{1,2}(0, T; W^{1,2}(\Omega; \mathbb{R}^n))$ as regularity of the deformation looks reasonable. This is a physically reasonable solution space: Since solids undergoing a phase transformation are likely to form microstructures, the regularity of the deformation will be low and one cannot necessarily expect the second spatial derivatives to be in $L^p(\Omega; \mathbb{R})$. Accepting this, we arrive at a heat equation in $L^1(\Omega; \mathbb{R})$. Since the heat capacity is constant, temperature is proportional to energy density, which is naturally measured in an $L^1(\Omega; \mathbb{R})$ norm. Therefore, the treatment of the heat equation in $L^1(\Omega; \mathbb{R})$ seems to be physically reasonable.

3.2. Statement of the Theorems

Now we are in a position to define a weak-renormalized solution of (6a)–(7c).




FIG. 2. Prototypical shape of $\phi(\theta)$, satisfying $\phi'(r_0) = 0$ for some point $r_0 > 0$ in addition to (E3). Here, ϕ is chosen to be constant close to the absolute temperature $\theta = 0$, which is plotted at the origin.

DEFINITION 3.2. A pair (u, θ) with $u: \Omega \times]0, T[\rightarrow \mathbb{R}^n$ and $\theta: \Omega \times]0, T[\rightarrow \mathbb{R}$ is said to be a *weak renormalized solution of the system (6a)–(7c)* if it satisfies the following conditions:

(i) Regularity of u :

$$\begin{aligned} u \in & L^\infty(W_g^{1,2}(\Omega; \mathbb{R}^n)) \cap W^{1,\infty}(0, T; L^2(\Omega; \mathbb{R}^n)) \\ & \cap W^{1,2}(0, T; W^{1,2}(\Omega; \mathbb{R}^n)) \cap W^{2,2}(0, T; W^{-1,2}(\Omega; \mathbb{R}^n)), \end{aligned}$$

(ii) u is a weak solution of (6a)–(6d): for every $\zeta \in C_0^\infty(\Omega \times]0, T[; \mathbb{R}^n)$,

$$\begin{aligned} \int_0^T \int_\Omega [(\sigma(\nabla u, \theta) + \nabla u_t) : \nabla \zeta - u_t \cdot \zeta_t] \, dx \, dt &= 0, \\ u(\cdot, 0) &= u_0(\cdot) \quad \text{in } \Omega \times \{0\}, \\ u_t(\cdot, 0) &= v_0(\cdot) \quad \text{in } \Omega \times \{0\}, \end{aligned}$$

(iii) θ is a renormalized solution of (7a)–(7c).

The main theorem of this paper can be formulated as follows.

THEOREM 3.1. *Let Ω be a nonempty, bounded domain in \mathbb{R}^n ($n = 2$ or $n = 3$) with a Lipschitz boundary. Assume the initial and boundary conditions satisfy the conditions stated in Section 2.1 and Φ satisfies hypotheses (E1)–(E7). Then there exists a weak renormalized solution of system (6a)–(7c).*

One also has the following result concerning the positivity of the temperature.

THEOREM 3.2. *Suppose in addition to the assumptions of Theorem 3.1 that $f(r_0) = 0$ for some $r_0 \geq 0$ and $\theta_0 \geq r_0$ almost everywhere in $\Omega \times]0, T[$.*

Then the temperature θ of the weak renormalized solution of system (6a)–(7c) satisfies $\theta \geq r_0$ almost everywhere in $\Omega \times]0, T[$.

We remark that Theorem 3.2 gives in particular for the prototypical energy given in Remark 2.2 the estimate $\theta \geq 0$ almost everywhere. To obtain $\theta > 0$, one has to modify the toy model for the energy to full-fill $\phi'(r_0) = 0$ for some point $r_0 > 0$. This can easily be achieved, an example of such a function is plotted in Figure 3.2.

4. PROOF OF THEOREMS 3.1 AND 3.2

The main part of this section will be devoted to the proof of Theorem 3.1. The proof of Theorem 3.2 follows at the end of the section; it is a straightforward application of ideas of Blanchard and Guibé [2].

We will use the Schauder Fixed Point Theorem to prove Theorem 3.1. Let $\hat{\theta} = \hat{\theta}(x, t)$ be an arbitrary element in $L^1(0, T; L^1(\Omega; \mathbb{R}))$. First, we consider the problem

$$u_{tt} = \text{Div} \left(\phi(\hat{\theta})\phi_1(\nabla u) + \phi_2(\nabla u) + \nabla u_t \right) \quad \text{in } \Omega \times]0, T[, \quad (9a)$$

$$u = g \quad \text{on } \partial\Omega \times [0, T[, \quad (9b)$$

$$u = u_0 \quad \text{in } \Omega \times \{0\}, \quad (9c)$$

$$u_t = v_0 \quad \text{in } \Omega \times \{0\}. \quad (9d)$$

We will show that this system has a unique solution \hat{u} . Substitution of $\hat{\theta}$ and \hat{u} in the nonlinear term on the right-hand side of Equation (7a) will give a solution $\hat{\theta}$ of (7a)–(7c). We will investigate continuity and compactness of the map $\Psi: \hat{\theta} \rightarrow \theta$.

THEOREM 4.1. *The system (9a)–(9d) admits a unique weak solution $\hat{u} = \hat{u}(x, t)$ with the regularity $\hat{u} \in L^\infty(W_g^{1,2}(\Omega; \mathbb{R}^n)) \cap W^{1,\infty}(0, T; L^2(\Omega; \mathbb{R}^n)) \cap W^{1,2}(0, T; W^{1,2}(\Omega; \mathbb{R}^n)) \cap W^{2,2}(0, T; W^{-1,2}(\Omega; \mathbb{R}^n))$.*

Proof. Since the temperature is fixed, this is a purely viscoelastic problem. Existence and uniqueness of a solution follows from the work of Friesecke and Dolzmann [14, Theorem 4.1]: a semi-implicit discretization in time leads to a variational problem. The integrand is convex, due to the discretized viscosity. Since the functional can be shown to be coercive (here, we need the assumption that $\partial\Omega$ is Lipschitz), a solution exists. Using different approximations in time, one can easily obtain most of the necessary weak convergences (to obtain strong convergence of the velocity, we use

again that $\partial\Omega$ is Lipschitz to apply an Aubin-type argument). The crucial step is to show strong convergence of ∇u ; this is [14, Proposition 3.1]. \square

A multiplication of equation (9a) by \hat{u}_t and integration over space and time yields

$$\begin{aligned} & \frac{1}{2} \int_{\Omega} |\hat{u}_t(t)|^2 dx + \int_0^t \int_{\Omega} |\nabla \hat{u}_t|^2 dx ds + \int_{\Omega} \Phi_2(\nabla \hat{u})(t) dx \\ &= - \int_0^t \int_{\Omega} \phi(\hat{\theta}) \phi_1(\nabla \hat{u}) : \nabla \hat{u}_t dx ds + \frac{1}{2} \int_{\Omega} |\hat{u}_t(0)|^2 dx + \int_{\Omega} \Phi_2(\nabla \hat{u})(0) dx \end{aligned}$$

for almost every t in $]0, T[$.

By (E7), there are positive constants c and c' such that

$$c |\nabla \hat{u}(t)|^2 - c' < \Phi_2(\nabla \hat{u})(t) \text{ and } \Phi_2(\nabla \hat{u})(0) < c' (|\nabla \hat{u}(0)|^2 + 1).$$

Hence, we obtain

$$\begin{aligned} & \frac{1}{2} \int_{\Omega} |\hat{u}_t(t)|^2 dx + \int_0^t \int_{\Omega} |\nabla \hat{u}_t|^2 dx ds + c \int_{\Omega} |\nabla \hat{u}(t)|^2 dx \quad (10) \\ & \leq - \int_0^t \int_{\Omega} \phi(\hat{\theta}) \phi_1(\nabla \hat{u}) : \nabla \hat{u}_t dx ds + \frac{1}{2} \|v_0\|_{L^2(\Omega; \mathbb{R}^n)}^2 \\ & \quad + c' \|u_0\|_{W^{1,2}(\Omega; \mathbb{R}^n)}^2 + C \end{aligned}$$

for almost every t in $]0, T[$.

By (E4), $\phi_1(\nabla \hat{u})$ is bounded. Using this and Young's inequality, the first term on the right-hand side can be estimated as follows:

$$\begin{aligned} & - \int_0^t \int_{\Omega} \phi(\hat{\theta}) \phi_1(\nabla \hat{u}) : \nabla \hat{u}_t dx ds \\ & \leq C \int_0^t \int_{\Omega} |\phi(\hat{\theta})|^2 dx ds + \frac{1}{2} \int_0^t \int_{\Omega} |\nabla \hat{u}_t|^2 dx ds. \quad (11) \end{aligned}$$

Combining (10) and (11), we arrive at

$$\begin{aligned} & \frac{1}{2} \int_{\Omega} |\hat{u}_t(t)|^2 dx + \frac{1}{2} \int_0^t \int_{\Omega} |\nabla \hat{u}_t|^2 dx ds + c \int_{\Omega} |\nabla \hat{u}(t)|^2 dx \\ & \leq C \int_0^t \int_{\Omega} |\phi(\hat{\theta})|^2 dx ds + \frac{1}{2} \|v_0\|_{L^2(\Omega; \mathbb{R}^n)}^2 + c' \|u_0\|_{W^{1,2}(\Omega; \mathbb{R}^n)}^2 + C \end{aligned}$$

for almost every t in $]0, T[$.

Since the terms on the left hand side are nonnegative, we can take the supremum over t . This yields

$$\begin{aligned} & \frac{1}{2} \|\hat{u}_t\|_{L^\infty(0,T;L^2(\Omega;\mathbb{R}^n))}^2 + \frac{1}{2} \|\nabla \hat{u}_t\|_{L^2(0,T;L^2(\Omega;\mathbb{R}^{n \times n}))}^2 \\ & \quad + c \|\nabla \hat{u}\|_{L^\infty(0,T;L^2(\Omega;\mathbb{R}^{n \times n}))}^2 \\ & \leq C \left[\|\phi(\hat{\theta})\|_{L^2(0,T;L^2(\Omega;\mathbb{R}))}^2 + \|v_0\|_{L^2(\Omega;\mathbb{R}^n)}^2 + \|u_0\|_{W^{1,2}(\Omega;\mathbb{R}^n)}^2 + 1 \right] \leq C, \end{aligned}$$

where C is independent of $\hat{\theta}$ by (E3). Poincaré's inequality

$$\|\hat{u} - g\|_{L^2(\Omega;\mathbb{R}^n)}^2 \leq C \left(\|\nabla \hat{u}\|_{L^2(\Omega;\mathbb{R}^{n \times n})}^2 + 1 \right)$$

finally gives the bound

$$\begin{aligned} & \frac{1}{2} \|\hat{u}_t\|_{L^\infty(0,T;L^2(\Omega;\mathbb{R}^n))}^2 \\ & \quad + \frac{1}{2} \|\nabla \hat{u}_t\|_{L^2(0,T;L^2(\Omega;\mathbb{R}^{n \times n}))}^2 + \|\hat{u}\|_{L^\infty(0,T;W^{1,2}(\Omega;\mathbb{R}^n))}^2 \leq C. \end{aligned}$$

Next, we prove continuous dependence of \hat{u} on $\hat{\theta}$. Let $\hat{\theta}_1, \hat{\theta}_2$ be two temperatures. Denote the corresponding solutions of (9a)–(9d) by \hat{u}_1 and \hat{u}_2 . A multiplication of the differences of the two equations by $\hat{u}_1 - \hat{u}_2$ and integration over space and time yields

$$\begin{aligned} & \partial_t \frac{1}{2} \left(\int_{\Omega} |\hat{u}_1(t) - \hat{u}_2(t)|^2 dx + \int_0^t \int_{\Omega} |\nabla \hat{u}_1 - \nabla \hat{u}_2|^2 dx ds \right) \quad (12) \\ & = - \int_0^t \int_{\Omega} \left[\left(\phi(\hat{\theta}_1) \phi_1(\nabla \hat{u}_1) - \phi(\hat{\theta}_2) \phi_1(\nabla \hat{u}_2) \right) : (\nabla \hat{u}_1 - \nabla \hat{u}_2) \right] dx ds \\ & \quad - \int_0^t \left[\int_{\Omega} (\phi_2(\nabla \hat{u}_1) - \phi_2(\nabla \hat{u}_2)) : (\nabla \hat{u}_1 - \nabla \hat{u}_2) \right] dx ds \\ & \quad + \int_0^t \int_{\Omega} |\partial_t \hat{u}_1 - \partial_t \hat{u}_2|^2 dx ds \\ & \leq \left| \int_0^t \int_{\Omega} \left(\phi(\hat{\theta}_1) - \phi(\hat{\theta}_2) \right) \phi_1(\nabla \hat{u}_1) : (\nabla \hat{u}_1 - \nabla \hat{u}_2) \right. \\ & \quad \left. + \phi(\hat{\theta}_2) (\phi_1(\nabla \hat{u}_1) - \phi_1(\nabla \hat{u}_2)) : (\nabla \hat{u}_1 - \nabla \hat{u}_2) dx ds \right| \\ & \quad + \text{Lip}(\phi_2) \int_0^t \int_{\Omega} |\nabla \hat{u}_1 - \nabla \hat{u}_2|^2 dx ds + \int_0^t \int_{\Omega} |\partial_t \hat{u}_1 - \partial_t \hat{u}_2|^2 dx ds \end{aligned}$$

$$\begin{aligned}
&\leq C \int_0^t \int_{\Omega} |\phi(\hat{\theta}_1) - \phi(\hat{\theta}_2)|^2 dx ds + \frac{1}{2} \int_0^t \int_{\Omega} |\nabla \hat{u}_1 - \nabla \hat{u}_2|^2 dx ds \\
&\quad + C \text{Lip}(\phi_1) \int_0^t \int_{\Omega} |\nabla \hat{u}_1 - \nabla \hat{u}_2|^2 dx ds \\
&\quad + \text{Lip}(\phi_2) \int_0^t \int_{\Omega} |\nabla \hat{u}_1 - \nabla \hat{u}_2|^2 dx ds + \int_0^t \int_{\Omega} |\partial_t \hat{u}_1 - \partial_t \hat{u}_2|^2 dx ds \\
&\leq C \int_0^t \int_{\Omega} |\phi(\hat{\theta}_1) - \phi(\hat{\theta}_2)|^2 dx ds + C \int_0^t \int_{\Omega} |\nabla \hat{u}_1 - \nabla \hat{u}_2|^2 dx ds \\
&\quad + \int_0^t \int_{\Omega} |\partial_t \hat{u}_1 - \partial_t \hat{u}_2|^2 dx ds
\end{aligned}$$

for almost every t in $]0, T[$ (the first inequality uses (E6), the second one Young's inequality, (E3), (E4) and (E5)). Similarly, using $\partial_t \hat{u}_1 - \partial_t \hat{u}_2$ as test function. and invoking (E3), (E4), (E5), (E6) and Young's inequality, one obtains

$$\begin{aligned}
&\partial_t \frac{1}{2} \int_0^t \int_{\Omega} |\partial_t \hat{u}_1 - \partial_t \hat{u}_2|^2 dx ds \tag{13} \\
&\leq \partial_t \frac{1}{2} \int_0^t \int_{\Omega} |\partial_t \hat{u}_1 - \partial_t \hat{u}_2|^2 dx ds + \frac{1}{2} \int_0^t \int_{\Omega} |\nabla \partial_t \hat{u}_1 - \nabla \partial_t \hat{u}_2|^2 dx ds \\
&= - \int_0^t \int_{\Omega} \left(\phi(\hat{\theta}_1) \phi_1(\nabla \hat{u}_1) - \phi(\hat{\theta}_2) \phi_1(\nabla \hat{u}_2) \right) : (\nabla \partial_t \hat{u}_1 - \nabla \partial_t \hat{u}_2) dx ds \\
&\quad - \int_0^t \int_{\Omega} \left(\phi_2(\nabla \hat{u}_1) - \phi_2(\nabla \hat{u}_2) \right) : (\nabla \partial_t \hat{u}_1 - \nabla \partial_t \hat{u}_2) dx ds \\
&\quad - \frac{1}{2} \int_0^t \int_{\Omega} |\nabla \partial_t \hat{u}_1 - \nabla \partial_t \hat{u}_2|^2 dx ds \\
&\leq C \int_0^t \int_{\Omega} |\phi(\hat{\theta}_1) - \phi(\hat{\theta}_2)|^2 dx ds + C \int_0^t \int_{\Omega} |\nabla \hat{u}_1 - \nabla \hat{u}_2|^2 dx ds
\end{aligned}$$

for almost every t in $]0, T[$.

Let us add (12) and (13). The second inequality uses $|\phi| \leq C$, the third one $|\phi'| \leq C$.

$$\begin{aligned}
&\partial_t \frac{1}{2} \left(\int_{\Omega} |\hat{u}_1(t) - \hat{u}_2(t)|^2 dx \tag{14} \right. \\
&\quad \left. + \int_0^t \int_{\Omega} |\nabla \hat{u}_1 - \nabla \hat{u}_2|^2 dx ds + \int_0^t \int_{\Omega} |\partial_t \hat{u}_1 - \partial_t \hat{u}_2|^2 dx ds \right) \\
&\leq C \int_0^t \int_{\Omega} |\phi(\hat{\theta}_1) - \phi(\hat{\theta}_2)|^2 dx ds
\end{aligned}$$

$$\begin{aligned}
 & + C \int_0^t \int_{\Omega} |\nabla \hat{u}_1 - \nabla \hat{u}_2|^2 \, dx \, ds + \int_0^t \int_{\Omega} |\partial_t \hat{u}_1 - \partial_t \hat{u}_2|^2 \, dx \, ds \\
 \leq & C \int_0^t \int_{\Omega} |\phi(\hat{\theta}_1) - \phi(\hat{\theta}_2)| \, dx \, ds \\
 & + C \int_0^t \int_{\Omega} |\nabla \hat{u}_1 - \nabla \hat{u}_2|^2 \, dx \, ds + \int_0^t \int_{\Omega} |\partial_t \hat{u}_1 - \partial_t \hat{u}_2|^2 \, dx \, ds \\
 \leq & C \int_0^t \int_{\Omega} |\hat{\theta}_1 - \hat{\theta}_2| \, dx \, ds \\
 & + C \int_0^t \int_{\Omega} |\nabla \hat{u}_1 - \nabla \hat{u}_2|^2 \, dx \, ds + \int_0^t \int_{\Omega} |\partial_t \hat{u}_1 - \partial_t \hat{u}_2|^2 \, dx \, ds
 \end{aligned}$$

for almost every t in $]0, T[$.

Hence, by Gronwall's inequality, \hat{u} depends continuously on $\hat{\theta}$. In particular, the map

$$\hat{\theta} \mapsto \nabla \hat{u}$$

is continuous from $L^1(0, T; L^1(\Omega; \mathbb{R}))$ to $L^2(0, T; L^2(\Omega; \mathbb{R}^{n \times n}))$; the second line of (13) shows that

$$\hat{\theta} \mapsto \nabla \hat{u}_t$$

is continuous from $L^1(0, T; L^1(\Omega; \mathbb{R}))$ to $L^2(0, T; L^2(\Omega; \mathbb{R}^n))$.

Using the estimates derived so far, one easily obtains a bound of the nonlinear term $f(\hat{\theta})\phi_1(\nabla \hat{u}) : \nabla \hat{u}_t + \nabla \hat{u}_t : \nabla \hat{u}_t$ on the right-hand side of the heat equation in $L^1(0, T; L^1(\Omega; \mathbb{R}))$. Namely, the following estimate uses (E3) to obtain boundedness of $f(\hat{\theta}) = \hat{\theta}\phi'(\hat{\theta})$:

$$\begin{aligned}
 & \left\| f(\hat{\theta})\phi_1(\nabla \hat{u}) : \nabla \hat{u}_t + \nabla \hat{u}_t : \nabla \hat{u}_t \right\|_{L^1(0, T; L^1(\Omega; \mathbb{R}))} \\
 & \leq C \left[\|u_0\|_{W^{1,2}(\Omega; \mathbb{R}^n)}^2 + \|v_0\|_{L^2(\Omega; \mathbb{R}^n)}^2 + 1 \right] \leq C,
 \end{aligned} \tag{15}$$

C being a constant independent of $\|\hat{\theta}\|_{L^1(0, T; L^1(\Omega; \mathbb{R}))}$.

Substituting \hat{u} and $\hat{\theta}$ in the right-hand side of the heat equation (7a), we get a unique solution θ :

$$\begin{aligned}
 \theta_t &= \Delta \theta + f(\hat{\theta})\phi_1(\nabla \hat{u}) : \nabla \hat{u}_t + \nabla \hat{u}_t : \nabla \hat{u}_t \quad \text{in } \Omega \times]0, T[, \\
 \theta &= 0 \quad \text{on } \partial\Omega \times]0, T[, \\
 \theta &= \theta_0 \quad \text{in } \Omega \times \{0\}
 \end{aligned} \tag{16}$$

(since the right hand side is in $L^1(0, T; L^1(\Omega; \mathbb{R}))$, it has a unique renormalized solution, see [2, Proposition 1 & 2]).

Recall that the map $\hat{\theta} \rightarrow \theta$ is denoted Ψ . The two maps $\hat{\theta} \mapsto \nabla \hat{u}_t$ and $\hat{\theta} \mapsto \nabla \hat{u}$ are continuous from $L^1(0, T; L^1(\Omega; \mathbb{R}))$ to $L^2(0, T; L^2(\Omega; \mathbb{R}^{n \times n}))$. Hence, the nonlinear term $f(\hat{\theta})\phi_1(\nabla \hat{u}) : \nabla \hat{u}_t + \nabla \hat{u}_t : \nabla \hat{u}_t$ of the right hand side of the heat equation is a continuous mapping of $L^1(0, T; L^1(\Omega; \mathbb{R}))$ to itself. Hence, by [2, Proposition 1], Ψ is continuous. Even more, by (15) and [2, Proposition 3], Ψ is compact for $1 \leq p < 1 + \frac{2}{n}$ as a map from $L^1(0, T; L^1(\Omega; \mathbb{R}))$ to $L^p(0, T; L^p(\Omega; \mathbb{R}))$.

The existence of a bounded set $B \subseteq L^1(0, T; L^1(\Omega; \mathbb{R}))$ with $\Psi(B) \subseteq B$ is now straightforward: inequality (15) gives a bound of the right-hand side of the heat equation by

$$C \left[\|u_0\|_{W^{1,2}(\Omega; \mathbb{R}^n)}^2 + \|v_0\|_{L^2(\Omega; \mathbb{R}^n)}^2 + 1 \right].$$

According to the theory of renormalized solutions (see, e.g., [2, Proposition 1]), there exists a constant C independent of $\|\hat{\theta}\|_{L^1(0, T; L^1(\Omega; \mathbb{R}))}$ such that

$$\|\theta\|_{L^1(0, T; L^1(\Omega; \mathbb{R}))} < R := C \left[\|u_0\|_{W^{1,2}(\Omega; \mathbb{R}^n)}^2 + \|v_0\|_{L^2(\Omega; \mathbb{R}^n)}^2 + \|\theta_0\|_{L^1(\Omega; \mathbb{R})} + 1 \right].$$

One can choose $B := B(0, R)$. An application of Schauder's Fixed Point Theorem finishes the proof. \square

We now sketch the proof of Theorem 3.2. It follows the proof of Theorem 4 in [2]. We denote by $\tilde{f}(r)$ the function

$$\tilde{f}(r) := \begin{cases} 0 & \text{if } r \leq r_0 \\ f(r) & \text{else} \end{cases}.$$

It is easily verified that the proof of Theorem 3.1 also applies to the system (6a)–(7c) with $f(\theta)$ replaced by $\tilde{f}(\theta)$. By considering for $K > 0$ the special test function $S(r) := \min(T_K(r - r_0), 0)$, one obtains from Definition 3.1 (iv) (χ_A is the characteristic function of a set A and $\tilde{S}(r) := \int_0^r S(x) dx$ the primitive of S)

$$\begin{aligned} & \int_{\Omega} \tilde{S}(\theta)(t) dx + \int_0^t \int_{\Omega} \chi_{]-K+r_0, r_0[} \nabla \theta^2 dx ds \\ &= \int_0^t \int_{\Omega} \left[\nabla u_t : \nabla u_t + \tilde{f}(\theta)\phi_1(\nabla u) : \nabla u_t \right] S(\theta) dx ds + \int_{\Omega} \tilde{S}(\theta_0) dx. \end{aligned}$$

By considering the signs of \tilde{f} and S , one obtains

$$\int_{\Omega} \tilde{S}(\theta)(t) dx \leq 0$$

for almost every t in $\Omega \times]0, T[$. This implies $\theta \geq r_0$ almost everywhere in $\Omega \times]0, T[$. The nature of the definition of \tilde{f} finally implies then

$$\tilde{f}(\theta) = f(\theta) \text{ almost everywhere in } \Omega \times]0, T[,$$

and Theorem 3.2 is established. \square

ACKNOWLEDGMENTS

This paper grew out of a part of the author's PhD thesis [27] written under the supervision of Prof. Jürgen Scheurle and Prof. Karl-Heinz Hoffmann. I thank both of them for support and advice. This research has been supported by the Deutsche Forschungsgemeinschaft, SFB 438. The article was written while the author held a postdoctoral position in the Division of Engineering and Applied Sciences, California Institute of Technology. Helpful comments of the referees are gratefully acknowledged.

REFERENCES

1. Kaushik Bhattacharya, Sergio Conti, Giovanni Zanzotto, and Johannes Zimmer. Symmetry and reversibility of martensitic transformations. Submitted to *Nature*, currently under revision.
2. D. Blanchard and O. Guibé. Existence of a solution for a nonlinear system in thermoviscoelasticity. *Adv. Differential Equations*, 5(10-12):1221–1252, 2000.
3. D. Blanchard and F. Murat. Renormalised solutions of nonlinear parabolic problems with L^1 data: existence and uniqueness. *Proc. Roy. Soc. Edinburgh Sect. A*, 127(6):1137–1152, 1997.
4. Dominique Blanchard. Truncations and monotonicity methods for parabolic equations. *Nonlinear Anal.*, 21(10):725–743, 1993.
5. Zhiming Chen and K.-H. Hoffmann. On a one-dimensional nonlinear thermoviscoelastic model for structural phase transitions in shape memory alloys. *J. Differential Equations*, 112(2):325–350, 1994.
6. C. M. Dafermos. Global smooth solutions to the initial-boundary value problem for the equations of one-dimensional nonlinear thermoviscoelasticity. *SIAM J. Math. Anal.*, 13(3):397–408, 1982.
7. C. M. Dafermos and L. Hsiao. Global smooth thermomechanical processes in one-dimensional nonlinear thermoviscoelasticity. *Nonlinear Anal.*, 6(5):435–454, 1982.
8. Andrea Dall'Aglio. Approximated solutions of equations with L^1 data. Application to the H -convergence of quasi-linear parabolic equations. *Ann. Mat. Pura Appl. (4)*, 170:207–240, 1996.
9. R. J. DiPerna and P.-L. Lions. On the Cauchy problem for Boltzmann equations: global existence and weak stability. *Ann. of Math. (2)*, 130(2):321–366, 1989.
10. R. J. DiPerna and P.-L. Lions. Ordinary differential equations, transport theory and Sobolev spaces. *Invent. Math.*, 98(3):511–547, 1989.
11. Patrick Dondl and Johannes Zimmer. Modeling and simulation of martensitic phase transitions with a triple point. Submitted to *Journal of the Mechanics and Physics of Solids*.

12. J. L. Ericksen. Weak Martensitic transformations in Bravais lattices. *Arch. Rational Mech. Anal.*, 107(1):23–36, 1989.
13. F. Falk. Landau theory and martensitic phase transitions. *J. Physique*, 43(NC-4):3–15, 1982.
14. G. Friesecke and G. Dolzmann. Implicit time discretization and global existence for a quasi-linear evolution equation with nonconvex energy. *SIAM J. Math. Anal.*, 28(2):363–380, 1997.
15. K.-H. Hoffmann and A. Żochowski. Analysis of the thermoelastic model of a plate with nonlinear shape memory alloy reinforcements. *Math. Methods Appl. Sci.*, 15(9):631–645, 1992.
16. Y. Huo and I. Müller. Interfacial and inhomogeneity penalties in phase transitions. To appear in *Contin. Mech. Thermodyn.*, 2003.
17. Song Jiang. Global large solutions to initial-boundary value problems in one-dimensional nonlinear thermoviscoelasticity. *Quart. Appl. Math.*, 51(4):731–744, 1993.
18. Pierre-Louis Lions. *Mathematical topics in fluid mechanics. Vol. 1.* The Clarendon Press Oxford University Press, New York, 1996. Incompressible models, Oxford Science Publications.
19. Irena Pawłow and Antoni Żochowski. Existence and uniqueness of solutions for a three-dimensional thermoelastic system. *Dissertationes Math. (Rozprawy Mat.)*, 406:46, 2002.
20. M. Pitteri. Reconciliation of local and global symmetries of crystals. *J. Elasticity*, 14(2):175–190, 1984.
21. Alain Prignet. Existence and uniqueness of “entropy” solutions of parabolic problems with L^1 data. *Nonlinear Anal.*, 28(12):1943–1954, 1997.
22. Reinhard Racke and Songmu Zheng. Global existence and asymptotic behavior in nonlinear thermoviscoelasticity. *J. Differential Equations*, 134(1):46–67, 1997.
23. Piotr Rybka. Dynamical modelling of phase transitions by means of viscoelasticity in many dimensions. *Proc. Roy. Soc. Edinburgh Sect. A*, 121(1-2):101–138, 1992.
24. Piotr Rybka. A priori estimates for the gradient of the solution to the system of viscoelasticity in several dimensions. *Topol. Methods Nonlinear Anal.*, 3(2):235–256, 1994.
25. Jürgen Sprekels. Global existence for thermomechanical processes with nonconvex free energies of Ginzburg-Landau form. *J. Math. Anal. Appl.*, 141(2):333–348, 1989.
26. Stephen J. Watson. Unique global solvability for initial-boundary value problems in one-dimensional nonlinear thermoviscoelasticity. *Arch. Ration. Mech. Anal.*, 153(1):1–37, 2000.
27. Johannes Zimmer. *Mathematische Modellierung und Analyse von Formgedächtnislegierungen in mehreren Raumdimensionen (Mathematical Modeling and Analysis of Shape Memory Alloys in Several Space Dimensions)*. PhD thesis, Technische Universität München, 2000.
28. Johannes Zimmer. Stored energy functions for phase transitions in crystals. Accepted for publication in *Archive for Rational Mechanics and Analysis*, 2003.