

**Max-Planck-Institut  
für Mathematik  
in den Naturwissenschaften  
Leipzig**

Necessary conditions for classifying  
m-separability of multipartite  
entanglements

by

*Wen Xu, Chuan-Jie Zhu, Zhu-Jun Zheng, and  
Shao-Ming Fei*

Preprint no.: 66

2020





# Necessary conditions for classifying **m**-separability of multipartite entanglements

Wen Xu<sup>1</sup>, Chuan-Jie Zhu<sup>2,3</sup>, Zhu-Jun Zheng<sup>1</sup> and Shao-Ming Fei<sup>4,5</sup>

<sup>1</sup> Department of Mathematics, South China University of Technology, GuangZhou 510640, P.R. China

<sup>2</sup> College of Mathematics and Physics Science,

Hunan University of Arts and Science, Changde 415000, P.R. China

<sup>3</sup> Department of Physics, Renmin University of China, Beijing 100872, P.R. China

<sup>4</sup> School of Mathematical Sciences, Capital Normal University, Beijing 100048, P.R.China

<sup>5</sup> Max-Planck-Institute for Mathematics in the Sciences, 04103 Leipzig, Germany

February 14, 2020

## Abstract

We study the norms of the Bloch vectors for arbitrary  $n$ -partite quantum states. A tight upper bound of the norms is derived for  $n$ -partite systems with different individual dimensions. These upper bounds are used to deal with the separability problems. Necessary conditions are presented for **m**-separable states in  $n$ -partite quantum systems. Based on the upper bounds, classification of multipartite entanglement is illustrated with detailed examples.

**Keywords** Bloch vectors · Norm · Upper bounds · Separability

## 1 Introduction

Quantum entanglement is a remarkable resource in the theory of quantum information, with numerous applications in quantum information processing, secure communication and channel protocols [1-3]. A multipartite quantum state that is not separable with respect to any bipartition is said to be genuinely multipartite entangled [4-6]. Genuinely multipartite entangled states have more advantages in quantum tasks compared with biseparable ones [7].

The notion of genuine multipartite entanglement (GME) was introduced in [7]. Let  $H_i^{d_i}, i = 1, \dots, n$ , denote  $d_i$ -dimensional Hilbert spaces. An  $n$ -partite state  $\rho \in H_1^{d_1} \otimes H_2^{d_2} \otimes \dots \otimes H_n^{d_n}$

$\cdots \otimes H_n^{d_n}$  can be expressed as  $\rho = \sum_i p_i |\psi_i\rangle\langle\psi_i|$ , where  $\sum_i p_i = 1$ ,  $0 < p_i \leq 1$ ,  $|\psi_i\rangle \in H_1^{d_1} \otimes H_2^{d_2} \otimes \cdots \otimes H_n^{d_n}$  are normalized pure states.  $\rho$  is biseparable if  $|\psi_i\rangle$  ( $i = 1, \dots, n$ ) can be expressed as one of the forms:  $|\psi_i\rangle = |\psi_i^{j_1 \dots j_{k-1}}\rangle \otimes |\psi_i^{j_k \dots j_n}\rangle$ , where  $|\psi_i^{j_1 \dots j_{k-1}}\rangle$  and  $|\psi_i^{j_k \dots j_n}\rangle$  denote pure states in  $H_i^{j_1} \otimes \cdots \otimes H_i^{j_{k-1}}$  and  $H_i^{j_k} \otimes \cdots \otimes H_i^{j_n}$ , respectively,  $j_1 \neq \cdots \neq j_n \in \{1, \dots, n\}$ ,  $k = 2, \dots, n-1$ . Otherwise,  $\rho$  is called genuine multipartite entangled. Correspondingly, we say that the state  $\rho = \sum_i p_i |\psi_i\rangle\langle\psi_i|$  is **m**-separable if all the  $|\psi_i\rangle$  are tensor products of m vectors in the subspaces of  $H_1^{d_1} \otimes H_2^{d_2} \otimes \cdots \otimes H_n^{d_n}$ .

Any quantum state has Bloch representation in multipartite high-dimensional quantum systems. By using the norms of the Bloch vectors, the density operators in lower dimensions were discussed in [8,9]. For bipartite and multipartite quantum states, separable conditions have been presented in [10-13]. The norms of the Bloch vectors for any qudit quantum states with subsystems less than or equal to four have been investigated in [14]. Two multipartite entanglement measures for n-qubit and n-qudit pure states are given in [15,16]. Some sufficient or necessary conditions of GME were presented in [17-19]. To the detection of GME, the common criterion is the entanglement witnesses [5,20-22]. In [6], the norms of the Bloch vectors give rise to a general framework to detect different classes of GME for arbitrary dimensional quantum systems.

In this paper, we study the Bloch representations of quantum states with arbitrary number of subsystems. In Section 2, we present tight upper bounds for the norms of Bloch vectors in n-qudit quantum states. These upper bounds are then used to derive tight upper bounds for entanglement measures in [15-16]. The upper bounds of the norms of the Bloch vectors are useful to study the separability. In Section 3, we investigate different subclasses of bi-separable states in n-partite systems. Necessary conditions for **m**-separability and complete classification of n-partite quantum systems are presented.

## 2 Upper bounds of the norms of Bloch vectors

Let  $\lambda_i$ ,  $i = 1, \dots, d^2 - 1$ , denote the generators of the special unitary group  $SU(d)$ , which satisfy  $\lambda_i^\dagger = \lambda_i$ ,  $Tr(\lambda_i) = 0$ ,  $Tr(\lambda_i \lambda_j) = 2\delta_{ij}$ . The following theorem gives the general result for n-partite quantum states.

**Theorem 1.** Let  $\rho \in H_1^{d_1} \otimes H_2^{d_2} \otimes \cdots \otimes H_n^{d_n}$  ( $n \geq 3$ ,  $2 \leq d_1 \leq d_2 \leq \cdots \leq d_n$ ) be an n-partite quantum state. We have

$$\|\mathbf{T}^{(12\dots n)}\|^2 \leq 2^n \left\{ 1 - \frac{\sum_{1 \leq i_1 < \dots < i_{n-1} \leq n} d_{i_1} \cdots d_{i_{n-1}} - 2d_n}{(n-2)d_1 \cdots d_{n-1} d_n^2} \right\}. \quad (1)$$

**Proof.**  $\rho$  has the Bloch representation:

$$\begin{aligned}
\rho = & \frac{1}{d_1 \cdots d_n} I_{d_1} \otimes \cdots \otimes I_{d_n} + \frac{1}{2} \left( \frac{1}{d_2 \cdots d_n} \sum_{i_1=1}^{d_1^2-1} t_{i_1}^{(1)} \lambda_{i_1} \otimes I_{d_2} \otimes \cdots \otimes I_{d_n} + \cdots \right. \\
& + \frac{1}{d_1 \cdots d_{n-1}} \sum_{i_n=1}^{d_n^2-1} t_{i_n}^{(n)} I_{d_1} \otimes I_{d_2} \otimes \cdots \otimes \lambda_{i_n} \Big) + \cdots \\
& + \frac{1}{2^n} \sum_{k=1}^n \sum_{i_k=1}^{d_k^2-1} t_{i_1 \cdots i_n}^{(1 \cdots n)} \lambda_{i_1} \otimes \lambda_{i_2} \otimes \cdots \otimes \lambda_{i_n},
\end{aligned} \tag{2}$$

where  $I_{d_i}$  denotes the  $d_i \times d_i$  identity matrix,  $i = 1, \dots, n$ ,  $t_{i_1}^{(1)} = \text{Tr}(\rho \lambda_{i_1} \otimes I_{d_2} \otimes \cdots \otimes I_{d_n})$ ,  $\dots$ ,  $t_{i_1 \cdots i_k}^{(j_1 \cdots j_k)} = \text{Tr}(\rho \lambda_{i_1} \otimes \cdots \otimes \lambda_{i_{j_k}} \otimes I_{d_{j_{k+1}}} \otimes \cdots \otimes I_{d_{j_n}})$ ,  $\dots$ ,  $t_{i_1 \cdots i_n}^{(1 \cdots n)} = \text{Tr}(\rho \lambda_{i_1} \otimes \lambda_{i_2} \otimes \cdots \otimes \lambda_{i_n})$  and  $\mathbf{T}^{(1)}, \dots, \mathbf{T}^{(j_1 \cdots j_k)}, \dots, \mathbf{T}^{(1 \cdots n)}$  are the vectors (tensors) with the elements  $t_{i_1}^{(1)}, \dots, t_{i_1 \cdots i_k}^{(j_1 \cdots j_k)}, \dots, t_{i_1 \cdots i_n}^{(1 \cdots n)}$  ( $1 \leq j_1 < \cdots < j_k \leq n$ ,  $i_s = 1, \dots, d_s^2 - 1$ ,  $s = 1, \dots, n$ ), respectively.

Set

$$\begin{aligned}
\|\mathbf{T}^{(1)}\|^2 &= \sum_{i_1=1}^{d_1^2-1} \left( t_{i_1}^{(1)} \right)^2, \\
&\quad \cdots, \\
\|\mathbf{T}^{(j_1 \cdots j_k)}\|^2 &= \sum_{s=1}^k \sum_{i_s=1}^{d_s^2-1} \left( t_{i_1 \cdots i_{j_k}}^{(j_1 \cdots j_k)} \right)^2, \\
&\quad \cdots, \\
\|\mathbf{T}^{(1 \cdots n)}\|^2 &= \sum_{s=1}^n \sum_{i_s=1}^{d_s^2-1} \left( t_{i_1 \cdots i_n}^{(1 \cdots n)} \right)^2,
\end{aligned}$$

$$x_1 = \frac{1}{d_2 \cdots d_n} \|\mathbf{T}^{(1)}\|^2 + \cdots + \frac{1}{d_1 \cdots d_{n-1}} \|\mathbf{T}^{(n)}\|^2,$$

$$\begin{aligned}
x_2 &= \frac{1}{d_3 \cdots d_n} \|\mathbf{T}^{(12)}\|^2 + \cdots + \frac{1}{d_1 \cdots d_{n-2}} \|\mathbf{T}^{(n-1,n)}\|^2, \\
&\quad \cdots, \\
x_n &= \|\mathbf{T}^{(12 \cdots n)}\|^2.
\end{aligned}$$

For a pure state  $\rho = |\psi\rangle\langle\psi|$ , one has  $\text{Tr}(\rho^2) = 1$ , namely,

$$\text{Tr}(\rho^2) = \frac{1}{d_1 \cdots d_n} + \frac{1}{2} x_1 + \frac{1}{2^2} x_2 + \cdots + \frac{1}{2^n} x_n = 1. \tag{3}$$

In the following we denote  $\rho_{j_1}, \rho_{j_2 \cdots j_n}$  the reduced density matrix for the subsystem  $H_{j_1}^{d_{j_1}}$  and  $H_{j_2}^{d_{j_2}} \otimes \cdots \otimes H_{j_n}^{d_{j_n}}$ ,  $j_1 \neq \cdots \neq j_n \in \{1, 2, \dots, n\}$ . One computes that,

$$\rho_{j_1} = \frac{1}{d_{j_1}} I_{d_{j_1}} + \frac{1}{2} \sum_{i_{j_1}=1}^{d_{j_1}^2-1} t_{i_{j_1}}^{(j_1)} \lambda_{i_{j_1}},$$

$$\begin{aligned}
\rho_{j_2 \cdots j_n} = & \frac{1}{d_{j_2} \cdots d_{j_n}} I_{d_{j_2}} \otimes \cdots \otimes I_{d_{j_n}} + \\
& \frac{1}{2} \left( \frac{1}{d_{j_3} \cdots d_{j_n}} \sum_{i_{j_3}=1}^{d_{j_3}^2-1} t_{i_{j_3}}^{(j_3)} \lambda_{i_{j_3}} \otimes \cdots \otimes I_{d_{j_n}} + \cdots \right. \\
& \left. + \frac{1}{d_{j_2} \cdots d_{j_{n-1}}} \sum_{i_{j_n}=1}^{d_{j_n}^2-1} t_{i_{j_n}}^{(j_n)} I_{d_{j_2}} \otimes \cdots \otimes \lambda_{i_{j_n}} \right) + \cdots \\
& + \frac{1}{2^{n-1}} \sum_{s=2}^n \sum_{i_{j_s}=1}^{d_{j_s}^2-1} t_{i_{j_2} \cdots i_{j_n}}^{(j_2 \cdots j_n)} \lambda_{i_{j_2}} \otimes \cdots \otimes \lambda_{i_{j_n}}.
\end{aligned}$$

For a pure state  $\rho = |\psi\rangle\langle\psi|$ , we have

$$Tr(\rho_{j_1}^2) = Tr(\rho_{j_2 \cdots j_n}^2), \quad (4)$$

which holds for any  $j_1 \neq \cdots \neq j_n \in \{1, 2, \dots, n\}$ . We obtain

$$\sum_{j_1=1}^n \frac{1}{d_{j_1}} Tr(\rho_{j_1}^2) = \sum_{j_1=1}^n \frac{1}{d_{j_1}} Tr(\rho_{j_2 \cdots j_n}^2). \quad (5)$$

Hence we get

$$\begin{aligned}
\sum_{i=1}^n \frac{1}{d_i^2} + \frac{1}{2} \sum_{i=1}^n \frac{1}{d_i^2} \|\mathbf{T}^{(i)}\|^2 = & \frac{n}{d_1 \cdots d_n} + \frac{n-1}{2} x_1 + \\
& \frac{n-2}{2^2} x_2 + \cdots + \frac{1}{2^{n-1}} x_{n-1}.
\end{aligned}$$

Then

$$\begin{aligned}
\frac{1}{2^2} x_2 = & \frac{1}{n-2} \left( \sum_{i=1}^n \frac{1}{d_i^2} - \frac{n}{d_1 \cdots d_n} \right) + \\
& \frac{1}{2(n-2)} \left( \sum_{i=1}^n \frac{1}{d_i} \|\mathbf{T}^{(i)}\|^2 - (n-1)x_1 \right) \\
& - \cdots - \frac{1}{2^{n-1}(n-2)} x_{n-1}.
\end{aligned} \quad (6)$$

Substituting (6) into (3), we get

$$\begin{aligned}
\frac{1}{2^n}x_n &= \left[ 1 - \frac{1}{n-2} \left( \sum_{i=1}^n \frac{1}{d_i^2} - \frac{2}{d_1 \cdots d_n} \right) \right] - \\
&\quad \frac{1}{2(n-2)} \left( \sum_{i=1}^n \frac{1}{d_i} \|\mathbf{T}^{(i)}\|^2 - x_1 \right) - \cdots - \frac{n-3}{2^{n-1}(n-2)} x_{n-1} \\
&\leq 1 - \frac{\sum_{1 \leq i_1 < \cdots < i_{n-1} \leq n} d_{i_1}^2 \cdots d_{i_{n-1}}^2 - 2d_1 \cdots d_n}{(n-2)d_1^2 \cdots d_n^2} \\
&\leq 1 - \frac{d_1 \cdots d_{n-1} \left( \sum_{1 \leq i_1 < \cdots < i_{n-1} \leq n} d_{i_1} \cdots d_{i_{n-1}} - 2d_n \right)}{(n-2)d_1^2 \cdots d_n^2} \\
&= 1 - \frac{\sum_{1 \leq i_1 < \cdots < i_{n-1} \leq n} d_{i_1} \cdots d_{i_{n-1}} - 2d_n}{(n-2)d_1 \cdots d_{n-1} d_n^2},
\end{aligned}$$

which is just

$$x_n = 2^n \left\{ 1 - \frac{\sum_{1 \leq i_1 < \cdots < i_{n-1} \leq n} d_{i_1} \cdots d_{i_{n-1}} - 2d_n}{(n-2)d_1 \cdots d_{n-1} d_n^2} \right\}, \quad n \geq 3.$$

Then we consider a mixed state  $\rho$  with ensemble representation  $\rho = \sum_i p_i |\psi_i\rangle\langle\psi_i|$ , where  $\sum_i p_i = 1$ ,  $0 < p_i \leq 1$ , by the convexity of the Frobenius norm one derives

$$\begin{aligned}
\|\mathbf{T}^{(12\cdots n)}(\rho)\|^2 &= \left\| \sum_i p_i \mathbf{T}^{(12\cdots n)}(|\psi_i\rangle\langle\psi_i|) \right\|^2 \\
&\leq \sum_i p_i \|\mathbf{T}^{(12\cdots n)}(|\psi_i\rangle\langle\psi_i|)\|^2 \\
&\leq 2^n \left\{ 1 - \frac{\sum_{1 \leq i_1 < \cdots < i_{n-1} \leq n} d_{i_1} \cdots d_{i_{n-1}} - 2d_n}{(n-2)d_1 \cdots d_{n-1} d_n^2} \right\}, \quad n \geq 3,
\end{aligned}$$

which ends the proof. ■

As a special case, consider  $d_1 = \cdots = d_n = d$  in Theorem 1. We have

**Corollary 1.** Let  $\rho \in H_1^d \otimes H_2^d \otimes \cdots \otimes H_n^d$  ( $n \geq 3, d \geq 2$ ) be an  $n$ -qudit quantum state. We have

$$\|\mathbf{T}^{(12\cdots n)}\|^2 \leq \frac{2^n [(n-2)d^n - nd^{n-2} + 2]}{(n-2)d^n}. \quad (7)$$

**Remark 1:** Corollary 1 is the generalization of the results of [14]. When  $n = 3, 4$ , the results of Corollary 1 reduce to the ones in [14].

The Bloch vectors are used to define a valid entanglement measure in [15, 16] as follows. For an  $n$ -qudit pure state, the entanglement measure is defined as:

$$E_{\mathbf{T}}(|\psi\rangle) = \left( \frac{d}{2} \right)^{\frac{n}{2}} \|\mathbf{T}^{(1\cdots n)}\| - \left( \frac{d(d-1)}{2} \right)^{\frac{n}{2}}, \quad (8)$$

where  $\mathbf{T}^{(1\cdots n)}$  is defined as a vector with elements  $t_{i_1\cdots i_n}^{(1\cdots n)} = \text{Tr}(\rho \lambda_{i_1} \otimes \lambda_{i_2} \otimes \cdots \otimes \lambda_{i_n})$ . Our results can give rise to an upper bound of the entanglement:

**Corollary 2.** *For any  $n$ -qudit pure state  $\rho \in H_1^d \otimes H_2^d \otimes \cdots \otimes H_n^d$  ( $n \geq 3, d \geq 2$ ), the entanglement measure has the upper bounds:*

$$E_{\mathbf{T}}(|\psi\rangle) \leq \left(\frac{d}{2}\right)^{\frac{n}{2}} \left[ \sqrt{d^n - \frac{n}{n-2}d^{n-2} + \frac{2}{n-2}} - (d-1)^{\frac{n}{2}} \right]. \quad (9)$$

### 3 The Necessary conditions for m-separable states

Now we study separability problems of  $n$ -partite quantum systems based on the upper bounds of the norms of Bloch vectors. Let  $\rho \in H_1^{d_1} \otimes H_2^{d_2} \otimes \cdots \otimes H_n^{d_n}$ ,  $n \geq 3$ ,  $2 \leq d_1 \leq d_2 \leq \cdots \leq d_n$ . If  $\rho$  can be written as  $\rho = \sum_i p_i |\psi_i\rangle\langle\psi_i|$ , where  $\sum_i p_i = 1$ ,  $0 < p_i \leq 1$ ,  $|\psi_i\rangle$  is one of the following sets:  $\{|\phi_{j_1}\rangle \otimes |\phi_{j_2\cdots j_n}\rangle\}, \dots, \{|\phi_1\rangle \otimes \cdots \otimes |\phi_n\rangle\}, \{|\phi_{j_1 j_2}\rangle \otimes |\phi_{j_3\cdots j_n}\rangle\}, \dots$ , where  $j_1 \neq \cdots \neq j_n \in \{1, \dots, n\}$ . Then  $\rho$  is called  $(1, n-1)$  separable,  $\dots, (\underbrace{1, \dots, 1}_n, 1)$  separable,  $(2, n-2)$  separable,  $\dots$ , respectively.

**Lemma 1.** *Let  $\rho_j \in H_j^{d_j}$  ( $j = 1, \dots, n, d_j \geq 2$ ) be the reduced density operator of  $\rho$ . We have*

$$\|\mathbf{T}^{(j)}\|^2 \leq \frac{2(d_j - 1)}{d_j}. \quad (10)$$

**Proof.**  $\rho_j$  has the Bloch representation:

$$\rho_j = \frac{1}{d_j} I_{d_j} + \frac{1}{2} \sum_{i_j=1}^{d_j^2-1} t_{i_j}^{(j)} \lambda_{i_j}, \quad (11)$$

where  $t_{i_j}^{(j)} = \text{Tr}(\rho_j \lambda_{i_j})$ ,  $\mathbf{T}^{(j)}$  is a vector with entries  $t_{i_j}^{(j)}$ ,  $j = 1, \dots, n$ ,  $i_j = 1, \dots, d_j^2 - 1$ . Since  $\text{Tr}(\rho_j^2) \leq 1$ , i.e.  $\frac{1}{d_j} + \frac{1}{2} \|\mathbf{T}^{(j)}\|^2 \leq 1$ , one obtains (10).  $\blacksquare$

**Lemma 2.** *Let  $\rho_{jk} \in H_j^{d_j} \otimes H_k^{d_k}$  ( $1 \leq j < k \leq n, 2 \leq d_j \leq d_k$ ) be the reduced density operator of  $\rho$ . We have*

$$\|\mathbf{T}^{(jk)}\|^2 \leq \frac{2^2(d_j^2 - 1)}{d_j^2}. \quad (12)$$

**Proof.**  $\rho_{jk}$  has the Bloch representation:

$$\begin{aligned} \rho_{jk} = & \frac{1}{d_j d_k} I_{d_j} \otimes I_{d_k} + \frac{1}{2d_k} \sum_{i_j=1}^{d_j^2-1} t_{i_j}^{(j)} \lambda_{i_j} \otimes I_{d_k} + \\ & \frac{1}{2d_j} \sum_{i_k=1}^{d_k^2-1} t_{i_k}^{(k)} I_{d_j} \otimes \lambda_{i_k} + \frac{1}{4} \sum_{i_j=1}^{d_j^2-1} \sum_{i_k=1}^{d_k^2-1} t_{i_j i_k}^{(jk)} \lambda_{i_j} \otimes \lambda_{i_k}, \end{aligned} \quad (13)$$

where  $t_{i_j}^{(j)} = \text{Tr}(\rho_{jk}\lambda_{i_j} \otimes I_{d_k})$ ,  $t_{i_k}^{(k)} = \text{Tr}(\rho_{jk}I_{d_j} \otimes \lambda_{i_k})$ ,  $t_{i_j i_k}^{(jk)} = \text{Tr}(\rho_{jk}\lambda_{i_j} \otimes \lambda_{i_k})$ .  $\mathbf{T}^{(j)}$ ,  $\mathbf{T}^{(k)}$  and  $\mathbf{T}^{(jk)}$  are vectors with entries  $t_{i_j}^{(j)}$ ,  $t_{i_k}^{(k)}$  and  $t_{i_j i_k}^{(jk)}$  ( $1 \leq j < k \leq n, i_j = 1, \dots, d_j^2 - 1, i_k = 1, \dots, d_k^2 - 1$ ). Set

$$\begin{aligned}\|\mathbf{T}^{(j)}\|^2 &= \sum_{i_j=1}^{d_j^2-1} \left(t_{i_j}^{(j)}\right)^2, \\ \|\mathbf{T}^{(k)}\|^2 &= \sum_{i_k=1}^{d_k^2-1} \left(t_{i_k}^{(k)}\right)^2, \\ \|\mathbf{T}^{(jk)}\|^2 &= \sum_{i_j=1}^{d_j^2-1} \sum_{i_k=1}^{d_k^2-1} \left(t_{i_j i_k}^{(jk)}\right)^2.\end{aligned}$$

For a pure state  $\rho_{jk} = |\psi\rangle\langle\psi|$ , one has  $\text{Tr}(\rho_{jk}^2) = 1$ , namely,

$$\text{Tr}(\rho_{jk}^2) = \frac{1}{d_j d_k} + \frac{1}{2d_k} \|\mathbf{T}^{(j)}\|^2 + \frac{1}{2d_j} \|\mathbf{T}^{(k)}\|^2 + \frac{1}{4} \|\mathbf{T}^{(jk)}\|^2 = 1. \quad (14)$$

Let  $\rho_j$  and  $\rho_k$  be the reduced density matrices with respect to the subsystems  $1 \leq j < k \leq n$ . Since for a pure state  $\rho_{jk}$ ,  $\text{Tr}(\rho_{jk}^2) = \text{Tr}(\rho_j^2)$ , i.e.  $\frac{1}{d_j} + \frac{1}{2} \|\mathbf{T}^{(j)}\|^2 = \frac{1}{d_k} + \frac{1}{2} \|\mathbf{T}^{(k)}\|^2$ . Therefore, we get

$$\begin{aligned}\|\mathbf{T}^{(jk)}\|^2 &= \frac{2^2(d_j^2 - 1)}{d_j^2} - \frac{2(d_j + d_k)}{d_j d_k} \|\mathbf{T}^{(j)}\|^2 \\ &\leq \frac{2^2(d_j^2 - 1)}{d_j^2}.\end{aligned} \quad (15)$$

Now we consider a mixed state  $\rho_{jk}$  with ensemble representation  $\rho_{jk} = \sum_i p_i |\psi_i\rangle\langle\psi_i|$ , where  $\sum_i p_i = 1$ ,  $0 < p_i \leq 1$ , by the convexity of the Frobenius norm one derives

$$\begin{aligned}\|\mathbf{T}^{(jk)}(\rho_{jk})\|^2 &= \left\| \sum_i p_i \mathbf{T}^{(jk)}(|\psi_i\rangle\langle\psi_i|) \right\|^2 \\ &\leq \sum_i p_i \|\mathbf{T}^{(jk)}(|\psi_i\rangle\langle\psi_i|)\|^2 \\ &\leq \frac{2^2(d_j^2 - 1)}{d_j^2},\end{aligned}$$

which ends the proof. ■

For the  $n$ -partite quantum state  $\rho \in H_1^{d_1} \otimes H_2^{d_2} \otimes \dots \otimes H_n^{d_n}$ ,  $n \geq 3, 2 \leq d_1 \leq \dots \leq d_n$ , consider the  $\mathbf{m}$ -partition of  $n$ -qudit quantum state  $\rho$ , we denote  $\mathbf{m} = (k_1, \dots, k_m)$ , where  $\sum_{s=1}^m k_s = n$ ,  $1 \leq k_1 \leq \dots \leq k_m \leq n - 1$  and  $n_j = \sum_{s=1}^j k_s$ ,  $1 \leq j \leq m, 1 \leq n_j \leq n$ . Moreover,

denotes the following:

$$\begin{aligned}
a_1 &= \frac{2(d_1-1)}{d_1}, \\
&\quad \cdots, \\
a_{n_p} &= \frac{2(d_{n_p}-1)}{d_{n_p}}, \\
a_{n_b} &= \frac{2^2(d_{n_b-1}^2-1)}{d_{n_b-1}^2}, \\
&\quad \cdots, \\
a_{n_{b+(q-1)}} &= \frac{2^2(d_{n_{b+(q-1)}-1}^2-1)}{d_{n_{b+(q-1)}-1}^2} \\
&\quad \cdots, \\
a_{n_c} &= 2^t \left\{ 1 - \frac{\sum d_{i_{n_c-(t-1)}} \cdots d_{i_{n_c-1}} - 2d_{n_c}}{(t-2)d_{n_c-(t-1)} \cdots d_{n_c-1} d_{n_c}^2} \right\}, \\
&\quad \cdots, \\
a_{n_{c+(s-1)}} &= 2^t \left\{ 1 - \frac{\sum d_{i_{n_{c+(s-1)}-(t-1)}} \cdots d_{i_{n_{c+(s-1)}-1}} - 2d_{n_{c+(s-1)}}}{(t-2)d_{n_{c+(s-1)}-(t-1)} \cdots d_{n_{c+(s-1)}-1} d_{n_{c+(s-1)}}^2} \right\}, \quad t \geq 3.
\end{aligned} \tag{16}$$

where  $p + q + \cdots + s = m$ ,  $p + 2q + \cdots + ts = n$ ,  $1 \leq p, b, q, \cdots, c, s \leq m$ .

Then from Lemmas 1, 2 and Theorem 1, we have

$$\begin{aligned}
\|\mathbf{T}^{(1)}\|^2 &\leq a_1, \\
&\quad \cdots, \\
\|\mathbf{T}^{(n_p)}\|^2 &\leq a_{n_p}, \\
\|\mathbf{T}^{(n_b-1, n_b)}\|^2 &\leq a_{n_b}, \\
&\quad \cdots, \\
\|\mathbf{T}^{(n_{b+(q-1)}-1, n_{b+(q-1)})}\|^2 &\leq a_{n_{b+(q-1)}}, \\
\|\mathbf{T}^{(n_c-(t-1), \cdots, n_c)}\|^2 &\leq a_{n_c}, \\
&\quad \cdots, \\
\|\mathbf{T}^{(n_{c+(s-1)}-(t-1), \cdots, n_{c+(s-1)})}\|^2 &\leq a_{n_{c+(s-1)}}, \quad t \geq 3.
\end{aligned}$$

where  $\mathbf{T}^{(u)}$  is a vector with the entries  $t_{i_u}^{(u)}$  ( $u = 1, \cdots, n_p$ ,  $i_u = 1, \cdots, d_u^2 - 1$ ).  $\mathbf{T}^{(xy)}$  is a vector with entries  $t_{i_x i_y}^{(xy)}$  ( $x = n_b - 1, \cdots, n_{b+(q-1)} - 1$ ,  $y = n_b, \cdots, n_{b+(q-1)}$ ,  $i_x = 1, \cdots, d_x^2 - 1$ ,  $i_y = 1, \cdots, d_y^2 - 1$ ).  $\cdots$ .  $\mathbf{T}^{(x_1, \cdots, x_s)}$  is a vector with entries  $t_{i_{x_1} \cdots i_{x_s}}^{(x_1 \cdots x_s)}$  ( $x_1 = n_c - (t-1), \cdots, n_{c+(s-1)} - (t-1)$ ,  $\cdots$ ,  $x_s = n_c, \cdots, n_{c+(s-1)}$ ,  $i_{x_1} = 1, \cdots, d_{x_1}^2 - 1, \cdots, i_{x_s} = 1, \cdots, d_{x_s}^2 - 1$ ).

The following theorem gives the necessary conditions of **m**-separability.

**Theorem 2.** Let  $\rho \in H_1^{d_1} \otimes H_2^{d_2} \otimes \cdots \otimes H_n^{d_n}$  ( $n \geq 3$ ,  $2 \leq d_1 \leq d_2 \leq \cdots \leq d_n$ ) be an  $n$ -partite quantum state. If  $\rho$  is **m**-separable we have

$$\|\mathbf{T}^{(12 \cdots n)}\|^2 \leq \prod_{f=1}^{n_p} a_f \prod_{g=n_b}^{n_{b+(q-1)}} a_g \cdots \prod_{h=n_c}^{n_{c+(s-1)}} a_h, \tag{17}$$

where  $\mathbf{m} = (k_1, \dots, k_m)$ ,  $\sum_{s=1}^m k_s = n$ ,  $1 \leq k_1 \leq \dots \leq k_m \leq n-1$  and  $n_j = \sum_{s=1}^j k_s$ ,  $1 \leq j \leq m$ ,  $1 \leq n_j \leq n$ ,  $a_f, a_g, a_h$  ( $f = 1, \dots, n_p$ ,  $g = n_b, \dots, n_{b+(q-1)}$ ,  $h = n_c, \dots, n_{c+(s-1)}$ ) are given in (16).

**Proof.** If  $\rho = |\psi\rangle\langle\psi|$  is an  $\mathbf{m}$ -separable pure state, where  $|\psi\rangle \in H_1^{d_1} \otimes H_2^{d_2} \otimes \dots \otimes H_n^{d_n}$ . Without lose of generality, assume that

$$|\psi\rangle = |\phi_1\rangle \otimes \dots \otimes |\phi_{n_p}\rangle \otimes |\phi_{n_b-1, n_b}\rangle \otimes \dots \otimes |\phi_{n_{b+(q-1)}-1, n_{b+(q-1)}}\rangle \otimes \dots \otimes |\phi_{n_c-(t-1), \dots, n_c}\rangle \otimes \dots \otimes |\phi_{n_{c+(s-1)}-(t-1), \dots, n_{c+(s-1)}}\rangle.$$

We have

$$\begin{aligned} t_{i_1 \dots i_n}^{(1 \dots n)} &= Tr(|\psi\rangle\langle\psi|\lambda_{i_1} \otimes \dots \otimes \lambda_{i_n}) \\ &= Tr(|\phi_1\rangle\langle\phi_1|\lambda_{i_1}) \dots Tr(|\phi_{n_p}\rangle\langle\phi_{n_p}|\lambda_{i_{n_p}}) \cdot \\ &\quad Tr\left(|\phi_{n_b-1, n_b}\rangle\langle\phi_{n_b-1, n_b}| \lambda_{i_{n_b-1}} \otimes \lambda_{i_{n_b}}\right) \dots \\ &\quad Tr\left(\left|\phi_{n_{b+(q-1)}-1, n_{b+(q-1)}}\right\rangle\left\langle\phi_{n_{b+(q-1)}-1, n_{b+(q-1)}}\right| \right. \\ &\quad \left.\lambda_{i_{n_{b+(q-1)}-1, n_{b+(q-1)}}} \otimes \lambda_{i_{n_{b+(q-1)}-1, n_{b+(q-1)}}}\right) \dots \\ &\quad Tr\left(\left|\phi_{n_c-(t-1), \dots, n_c}\right\rangle\left\langle\phi_{n_c-(t-1), \dots, n_c}\right| \right. \\ &\quad \left.\lambda_{i_{n_c-(t-1)}} \otimes \dots \otimes \lambda_{i_{n_c}}\right) \dots \\ &\quad Tr\left(\left|\phi_{n_{c+(s-1)}-(t-1), \dots, n_{c+(s-1)}}\right\rangle\left\langle\phi_{n_{c+(s-1)}-(t-1), \dots, n_{c+(s-1)}}\right| \right. \\ &\quad \left.\lambda_{i_{n_{c+(s-1)}-(t-1)}} \otimes \dots \otimes \lambda_{i_{n_{c+(s-1)}}}\right) \\ &= t_{i_1}^{(1)} \dots t_{i_{n_p}}^{(n_p)} t_{i_{n_b-1} i_{n_b}}^{(n_b-1, n_b)} \dots t_{i_{n_{b+(q-1)}-1} i_{n_{b+(q-1)}}}^{(n_{b+(q-1)}-1, n_{b+(q-1)})} \dots \\ &\quad t_{i_{n_c-(t-1)} \dots i_{n_c}}^{(n_c-(t-1), \dots, n_c)} \dots t_{i_{n_{c+(s-1)}-(t-1)} \dots i_{n_{c+(s-1)}}}^{(n_{c+(s-1)}-(t-1), \dots, n_{c+(s-1)})}. \end{aligned} \tag{18}$$

Thus

$$\begin{aligned} \|\mathbf{T}^{(12 \dots n)}\|^2 &= \|\mathbf{T}^{(1)}\|^2 \dots \|\mathbf{T}^{(n_p)}\|^2 \\ &\quad \|\mathbf{T}^{(n_b-1, n_b)}\|^2 \dots \|\mathbf{T}^{(n_{b+(q-1)}-1, n_{b+(q-1)})}\|^2 \dots \\ &\quad \|\mathbf{T}^{(n_c-(t-1), \dots, n_c)}\|^2 \dots \|\mathbf{T}^{(n_{c+(s-1)}-(t-1), \dots, n_{c+(s-1)})}\|^2 \\ &\leq \prod_{f=1}^{n_p} a_f \prod_{g=n_b}^{n_{b+(q-1)}} a_g \dots \prod_{h=n_c}^{n_{c+(s-1)}} a_h. \end{aligned} \tag{19}$$

Then for any mixed state  $\rho = \sum_k p_k |\psi_k\rangle\langle\psi_k| \in H_1^{d_1} \otimes H_2^{d_2} \otimes \cdots \otimes H_n^{d_n}$ , where  $\sum_k p_k = 1$ ,  $0 < p_k \leq 1$ , by the convexity of the Frobenius norm one derives

$$\begin{aligned} \|\mathbf{T}^{(12\cdots n)}(\rho)\|^2 &= \left\| \sum_k p_k \mathbf{T}^{(12\cdots n)}(|\psi_k\rangle\langle\psi_k|) \right\|^2 \\ &\leq \sum_k p_k \left\| \mathbf{T}^{(12\cdots n)}(|\psi_k\rangle\langle\psi_k|) \right\|^2 \\ &\leq \prod_{f=1}^{n_p} a_f \prod_{g=n_b}^{n_{b+(q-1)}} a_g \cdots \prod_{h=n_c}^{n_{c+(s-1)}} a_h. \end{aligned} \quad (20)$$

■

**Remark 2:** Set  $d_1 = \cdots = d_n = d$ . Then  $a_1 = \cdots = a_{n_p}$ ,  $a_{n_b} = \cdots = a_{n_{b+(q-1)}} = a_2, \dots$ ,  $a_{n_c} = \cdots = a_{n_{c+(s-1)}} = a_t$ . Then (17) gives rise to

$$\|\mathbf{T}^{(12\cdots n)}\|^2 \leq a_1^p a_2^q \cdots a_t^s. \quad (21)$$

**Remark 3:** Let  $\rho \in H_1^{d_1} \otimes H_2^{d_2} \otimes H_3^{d_3} \otimes H_4^{d_4}$  be a four-partite quantum state. One has

$$\|\mathbf{T}^{(1234)}\|^2 \leq \begin{cases} \frac{2^4(d_1-1)}{d_1} \left\{ 1 - \frac{d_2d_3+d_2d_4+d_3d_4-2d_4}{d_2d_3d_4^2} \right\}, & \text{if } \rho \text{ is (1, 3) separable;} \\ \frac{2^4(d_1^2-1)(d_3^2-1)}{d_1^2d_3^2}, & \text{if } \rho \text{ is (2, 2) separable;} \\ \frac{2^4(d_1-1)(d_2-1)(d_3^2-1)}{d_1d_2d_3^2}, & \text{if } \rho \text{ is (1, 1, 2) separable;} \\ \frac{2^4 \prod_{i=1}^4 (d_i-1)}{\prod_{i=1}^4 d_i}, & \text{if } \rho \text{ is (1, 1, 1, 1) separable.} \end{cases} \quad (22)$$

The following two examples show that the upper bounds in Theorem 2 are nontrivial and are tight.

*Example 1:* Consider the quantum state  $\rho \in H_1^2 \otimes H_2^2 \otimes H_3^2 \otimes H_4^2 \otimes H_5^2$ ,

$$\rho = x(|\psi\rangle\langle\psi| + |\varphi\rangle\langle\varphi|) + \frac{1-2x}{32} I_{32}, \quad (23)$$

where  $|\psi\rangle = \frac{1}{\sqrt{2}}(|00000\rangle + |11111\rangle)$ ,  $|\varphi\rangle = \frac{1}{2}(|00001\rangle + |00010\rangle + |00100\rangle + |01000\rangle)$ . Since  $\|\mathbf{T}^{(12345)}\|^2 = \sum_{i_1, \dots, i_5=1}^3 \left( t_{i_1 \dots i_5}^{(1 \dots 5)} \right)^2$ , where  $t_{i_1 \dots i_5}^{(1 \dots 5)} = \text{Tr}(\rho \lambda_{i_1}^{(1)} \otimes \dots \otimes \lambda_{i_5}^{(5)})$  are the entries of  $\mathbf{T}^{(12345)}$ , we have  $\|\mathbf{T}^{(12345)}\|^2 = 20x^2$ . Thus for  $\frac{3\sqrt{5}}{10} < x \leq \frac{\sqrt{15}}{5}$ ,  $\rho$  is not (1, 4) or (1, 2, 2) separable. For  $\frac{\sqrt{15}}{5} < x \leq 1$ ,  $\rho$  is not (2, 3) separable. For  $\frac{\sqrt{5}}{5} < x \leq \frac{3\sqrt{5}}{10}$ ,  $\rho$  is not (1, 1, 3) separable. For  $\frac{\sqrt{15}}{10} < x \leq \frac{\sqrt{5}}{5}$ ,  $\rho$  is not (1, 1, 1, 2) separable. For  $\frac{\sqrt{5}}{10} < x \leq \frac{\sqrt{15}}{10}$ ,  $\rho$  is not (1, 1, 1, 1, 1) separable.

*Example 2:* Consider the quantum state  $\rho \in H_1^2 \otimes H_2^3 \otimes H_3^4 \otimes H_4^5$ ,

$$\rho = x|\psi\rangle\langle\psi| + \frac{1-x}{120}I_{120}, \quad (24)$$

where  $|\psi\rangle = \frac{1}{\sqrt{2}}(|0\rangle_1|0\rangle_2|0\rangle_3|4\rangle_4 + |1\rangle_1|0\rangle_2|0\rangle_3|0\rangle_4)$ ,  $|0\rangle_1 := [1, 0]^T$ ,  $|1\rangle_1 := [0, 1]^T$ ,  $|0\rangle_2 := [1, 0, 0]^T$ ,  $|0\rangle_3 := [1, 0, 0, 0]^T$ ,  $|0\rangle_4 := [1, 0, 0, 0, 0]^T$ ,  $|4\rangle_4 := [0, 0, 0, 0, 1]^T$  ( $T$  is the transpose). Since  $\|\mathbf{T}^{(1234)}\|^2 = \sum_{k=1}^4 \sum_{i_k=1}^{d_k^2-1} \left(t_{i_1 \dots i_4}^{(1 \dots 4)}\right)^2$ , where  $t_{i_1 \dots i_4}^{(1 \dots 4)} = \text{Tr}(\rho \lambda_{i_1} \otimes \dots \otimes \lambda_{i_4})$  are the entries of  $\mathbf{T}^{(1234)}$ , we can compute that  $\|\mathbf{T}^{(1234)}\|^2 = 6x^2$ . Thus for  $\frac{\sqrt{263}}{15} < x \leq \frac{\sqrt{30}}{4}$ ,  $\rho$  is not  $(1, 3)$  separable. For  $\frac{\sqrt{30}}{4} < x \leq 1$ ,  $\rho$  is not  $(2, 2)$  separable. For  $\frac{\sqrt{30}}{6} < x \leq \frac{\sqrt{263}}{15}$ ,  $\rho$  is not  $(1, 1, 2)$  separable. For  $\frac{2\sqrt{30}}{15} < x \leq \frac{\sqrt{30}}{6}$ ,  $\rho$  is not  $(1, 1, 1, 1)$  separable.

From the above results, we are able to classify the entanglement of n-partite quantum states by using the norms of the Bloch vector  $\|\mathbf{T}^{(12 \dots n)}\|^2$ . The upper bounds of  $\|\mathbf{T}^{(12 \dots n)}\|^2$  can be used to identify the  $\mathbf{m}$ -separable n-partite quantum states, which include the fully separable states and the genuine multipartite entangled states as special classes.

## 4 Conclusion

Classification and detection of quantum entanglement are basic and fundamental problems in theory of quantum entanglement. We have investigated the norms of the Bloch vectors for arbitrary n-partite quantum systems. Tight upper bounds of the norms have been derived, and used to derive tight upper bounds for entanglement measure defined by the norms of Bloch vectors. The upper bounds have a close relationship to the separability. Necessary conditions have been presented for  $\mathbf{m}$ -separable quantum states. With these upper bounds a complete classification of n-partite quantum states has been obtained. Our results may highlight further studies on the quantum entanglement.

## Acknowledgments

This work is supported by the NSF of China under Grant Nos. 11571119 and 11675113, the Key Project of Beijing Municipal Commission of Education (Grant No. KZ201810028042), and Beijing Natural Science Foundation (Z190005). It is a pleasure to thank Jin-Wei Huang for helpful discussion.

## References

- [1] A. K. Ekert, Phys. Rev. Lett. 67, 661 (1991).
- [2] C. H. Bennett and S. J. Wiesner, Phys. Rev. Lett. 69, 2881 (1992).

- [3] C. H. Bennett, G. Brassard, C. Crepeau, R. Jozsa, A. Peres and W. K. Wootters, Phys. Rev.Lett. 70, 1895 (1993).
- [4] W. Dür, J. I. Cirac, and R. Tarrach, Phys. Rev. Lett. 83, 3562(1999).
- [5] M. Huber and R. Sengupta, Phys. Rev. Lett. 113, 100501 (2014).
- [6] J. I. de Vicente and M. Huber, Phys. Rev. A 84, 062306 (2011).
- [7] R. Horodecki, P. Horodecki, M. Horodecki, and K. Horodecki, Rev. Mod. Phys. 81, 865 (2009).
- [8] L. Jakbczyk, M. Siennicki, Phys. Lett. A 286, 383(2001).
- [9] G. Kimura, Phys. Lett. A 314, 339 (2003).
- [10] J. D. Vicente, Quantum Inf. Comput. 7, 624(2007).
- [11] J. D. Vicente, J. Phys. A: Math. and Theor., 41, 065309(2008).
- [12] A.S. M. Hassan, P. S. Joag, Quantum Inf. Comput. 8, 0773(2008).
- [13] M. Li, J. Wang, S.M. Fei and X.Q. Li-Jost, Phys. Rev. A, 89,022325(2014).
- [14] M. Li, Z.Wang, J.Wang, S.Q.Shen and S.M.Fei, Europhys. Lett. A 125, 20006 (2019).
- [15] A. S. M. Hassan, P. S. Joag, Phys. Rev. A, 77, 062334 (2008).
- [16] A. S. M. Hassan, P. S. Joag, Phys. Rev. A, 80, 042302 (2009).
- [17] P. van Loock and A. Furusawa, Phys. Rev. A 67, 052315 (2003).
- [18] M. J. Zhao, T. G. Zhang, X. Li-Jost and S. M. Fei, Phys. Rev. A 87, 012316 (2013).
- [19] M. Li, J. Wang, S. Shen, Z. Chen and S. M. Fei, Scientific Reports 7, 17274 (2017).
- [20] J. D. Bancal, N. Gisin, Y. C. Liang and S. Pironio, Phys. Rev. Lett. 106, 250404 (2011).
- [21] B. Jungnitsch, T. Moroder and O. Guhne, Phys. Rev. A 84, 032310 (2011).
- [22] J. Y. Wu, H. Kampermann, D. Bru, C. Klockl and M. Huber, Phys. Rev. A 86, 022319 (2012).